

Standard Procedure for the study of *Posidonia oceanica* Leaf Litter

Christine PERGENT-MARTINI\*, Gérard PERGENT\* and Javier ROMERO\*\*

\*LBMEB, Faculté des Sciences de Luminy, 13288 Marseille Cédex 09 (France)

\*\*Departamento de Ecología, Facultad de Biología, Universidad de Barcelona, 08028 Barcelona (España)

While leaf litter is recognized as one of the key compartments in the dynamics of terrestrial ecosystems (OLSON, 1963; KARKANIS, 1975), for the *Posidonia oceanica* bed it has only been studied sporadically (WITTMANN *et al.*, 1981; FRANCOUR, 1990).

As part of a general survey of the functioning of the *Posidonia oceanica* ecosystem ("A functional approach to the *Posidonia oceanica* ecosystem of the Mediterranean"), a standard procedure has been worked out for determining the structure, chemical composition and fragmentation and degradation mechanisms of *Posidonia* leaf litter.

Sampling is carried out by scuba diving, using a suction device. A quadrat 35 cm square is set up in a homogeneous area that is representative of the *Posidonia* bed under investigation. Within this quadrat, the living leaves are cut off at a height of 3 to 5 cm from the base and removed, and the shoots are counted. The leaf litter is collected in bags (1 mm mesh). Sampling is repeated three times for each station. The samples are transported from the sampling site to the laboratory in damp medium.

In each sample, non-litter elements (eg. living leaves, shell debris, algae, animals etc) are removed by hand. Dead rhizomes, with or without scales, and whole scales are separated from the litter, and constitute the RHIZOME FRACTION. Sorting by gravity is carried out to separate the litter from the sediment, which is discarded. Two sieves of different mesh size are used to sort the litter into a COARSE FRACTION (leaves larger than 8 mm) and a FINE FRACTION (leaf debris of between 1 mm and 8 mm).

The three fractions obtained for each sample are placed in the dryer at a temperature of 70°C, until a constant weight is obtained (usually 48 hours). After drying, the samples are weighed on precision scales (mg).

Leaf fragmentation experiments are carried out in situ. Aged adult leaves, that are still in place in the meadow, are collected and brought back to the laboratory. They are weighed (sub-samples of 30 +/- 0.5 g) and placed in bags (1 mm mesh) sealed with strips of Velcro. The bags are returned to the environment beneath the leaf cover in series of three. The series of bags are collected after a predetermined period of time (1, 2, 3, 4, 8, 12, 16 and 24 weeks). The samples collected are separated into three size classes: leaves with a length of > 5 cm (LARGE LEAVES); leaves with a length of between 5 cm and 8 mm (BROKEN LEAVES), and leaf debris of between 8 mm and 1 mm (DEBRIS). Each fraction is dried in the dryer at 70°C (constant weight), then weighed.

The totality of the samples is powdered (pulveriser) and sieved (0.63 µm mesh). The CHN content (Micro HN Determinator, CHN 800), the percentage of ash (Thermolyne Sybron type 2000, 8 hours at 550°C) and the Phosphorus content (Induction Coupling Plasma, after acid digestion according to DELGADO, 1986) are measured for each fraction.

Our preliminary results (ROMERO *et al.*, sous-pressé) would appear to suggest that the distribution patterns of leaf litter are subject to considerable variation according to the depth, time and site of sampling. At Ischia (Italy), leaf litter maxima (coarse fraction + fine fraction) are found at intermediate depths (Figure 1).




Figure 1: Mean litter biomass (in mg dry weight per quadrat) at Ischia (Italy), at various depths according to sampling date.

An investigation of degradation in situ, at -5 m and -20 m, has shown that depth does not appear to have any influence on the rate of degradation. On the other hand, the rate of degradation does depend on the month of the year of the investigation (higher degradation rate in July). The curve of decline is of the exponential type:

$$y = \exp(-0.0066x + 1.63) \quad -5 \text{ m - October 1988 experiment.}$$

$$y = \exp(-0.0087x + 1.78) \quad -20 \text{ m - July 1988 experiment.}$$

**Acknowledgements:** This study was made possible by financial support from the European Communities Commission and valuable assistance from the staff of the Ischia laboratory (Italy).

## References

- DELGADO, O., 1986. - Contenido de fósforo de los tejidos de fanerógamas marinas del Mediterráneo y su relación con la dinámica de cada especie. *Oecologia Aquatica*, 8: 139-151.
- FRANCOUR, P., 1990. - *Dynamique de l'écosystème à Posidonia oceanica dans le Parc National de Port-Cros. Analyse des compartiments matie, litière, faune vagile, échinodermes et poissons*. Thèse doctorat, Université Paris VI, 280 p.
- MARKANIS, M., 1975. - Decomposition of litter of various species of deciduous trees and its effect in soil environment. *Fragmenta Floristica et Geobotanica*, 21: 71-97.
- SAN, J.S., 1963. - Energy storage and the balance of producers and decomposers in ecological systems. *Ecology*, 44(2): 322-331.
- ROMERO, J., PERGENT, G., PERGENT-MARTINI, C., MATEO, M.A., REGNIER, C., sous presse. - The detritic compartment in a *Posidonia oceanica* meadow: litter features, decomposition rates and mineral stocks. *Marine Ecology*.
- WITTMANN, K., SCIPIONE M.B., FRESI E., 1981. - Some laboratory experiments on the activity of the macrofauna in the fragmentation of detrital leaves of *Posidonia oceanica* (L.) Delile. *Rapp. P.V. Réun. Comités Internation. Explor. sci. Médit.*, 27(2): 205-206.

The Regression of *Posidonia oceanica* Meadows in El Campello (Spain)

J.-L. SANCHEZ LIZASO, J.-E. GUILLEN NIETO and A.-A. RAMOS ESPLA\*

Institut d'Ecologia Litoral, Ctra. Benimagrell s/n. 03560 El Campello. Alicante (España)

## ABSTRACT

The regression of *Posidonia oceanica* meadows has been studied along a 7 km coastal sector on the El Campello littoral. Changes of shallow *Posidonia* beds in the last 30 years are described. Deep *Posidonia* meadow is badly damaged due to illegal trawling. The first symptoms of trawling are detected at 13 m depth. But the deeper we go, the more degraded the meadow gets, reaching densities under 1 sh/m<sup>2</sup>. Dead *Posidonia* is seen even suggest at 29 m depth. In order to protect the deep meadow, we suggest as a feasible solution the installation of artificial reefs.

## INTRODUCTION

*Posidonia oceanica* meadow regression has been studied by a great number of scientists (see PERES, 1984). Studies on the effects of illegal trawling in *Posidonia* beds have been carried out in the Tyrrhenian Sea (ARDIZZONE & MIGLIUOLO, 1982; ARDIZZONE & PELUSI, 1983, 1984).

If we concentrate on Spain's situation, we can state that trawling has been forbidden since 1962 at less than 50 m for the whole year (Orden de 7 de julio de 1962, Reglamento de la pesca de arrastre a remolque. B.O.E. N° 16). In summer trawling is even forbidden at less than 130m depth (Orden de 30 de julio de 1975 sobre pesca de arrastre en el Mediterraneo. B.O.E. N° 193). However, trawling ships often work on *P. oceanica* meadow at less depth.

## MATERIAL AND WORKING METHODOLOGY

This piece of research was conducted on a 7 km coastal sector at El Campello (Alicante, SE of Spain). In order to complete this study we carried out nine perpendicular transects to the coastline from the upper level of the *P. oceanica* meadows to a depth of 29m and, in addition, several precise dives. The obtained points were positioned by means of a sextant and enfilades to the coast (RAMOS, 1984). The reconstruction of the *Posidonia*'s upper level was made with the aid of aerial photographs taken in the years 1956, 1978 and 1987.

## RESULTS

Two little breakwaters have been built in the area of research, between 1956 and 1987. If we compare the aerial photographs, we can clearly see that shallow *Posidonia* meadows have moved back. The greatest regression appears in the north of Cala Baeza and La Coveta, between 1978 and 1987. At the same period the little port of Cala Baeza was filled up, and now boats cannot tie up in this port.

Deep *Posidonia oceanica* meadow is destroyed due to illegal otter trawling. The depth at which the first trawling symptoms appear increases gradually from north to south. In the northern part of the studied area, which is off Carritxal beach, the first degradation symptoms were observed at 13 m depth; in the intermediate sector, at Cala Baeza, they appear at either 15 or 16 m; finally, in the southern part, Barranc d'Aigues-Morro Blanc, at 17 m depth. At this depth 0.5 m wide channels parallel to the coast line are detected as well as pulled up rhizomes on account of the mechanical effect of trawl boards.

As we go deeper, channels become more frequent and wider; there seem to be more pulled up rhizomes and the proportion of dead matte increase. *Posidonia* meadow density decreases quickly until it reaches values under 1 sh/m<sup>2</sup>. From 22 m depth, the sea bottom presents a desolating sight; there are very few *Posidonia* spots and a great mass of dead matte frequently covered with light layers of sediments. *Posidonia oceanica* remains have been detected at 25 m depth at Carritxal and even at 29 m depth between Barranc d'Aigues and Morro Blanc.

## CONCLUSIONS

Regression in a shallow *Posidonia oceanica* meadow due to coastal line modifications is seen. More serious seems to be the regression due to illegal trawling.

It is believed that along 7 km of coast, 290 Ha of *Posidonia* meadow have been destroyed due to otter trawling. Unfortunately all the province of Alicante seems to be threatened by the same devastating problem.

Moreover several unpublished issues point out that the *Posidonia* meadow at Tabarca's island, La Vila Joiosa and El Campello is largely altered or destroyed at a depth which may fluctuate between 13 and 24 m. Likewise, according to some fishermen, the same thing is happening at other ports in the province of Alicante.

Vigilance committees have clearly proved to be inadequate and insufficient in the battle against illegal trawling and its ravaging effect on *Posidonia oceanica*. In our view, the only feasible solution to this problem is to install artificial antitrawling reefs, like those installed in Tabarca's Marine Reserve (Ramos *et al.*, in press).

## REFERENCES

- ARDIZZONE, G.D. & MIGLIUOLO, A. 1982. Modificazioni di una prateria di *Posidonia oceanica* (L.) Delile del Medio Tirreno sottoposta ad attività di pesca a strascico. Atti XIII Congr. Soc. Ital. Biol. Mar., Cefalù, It. su *Nat. sicil.*, S.IV, VI (Suppl.): 509-515.
- ARDIZZONE, G.D. & PELUSI, P. 1983. Fish population exposed to coastal bottom trawling along a Tyrrhenian Sea. *Rapp. Comités Internation. Explor. sci. Mer Médit.*, Monaco: 107-110.
- ARDIZZONE, G.D. & PELUSI, P. 1984. Yield and damage of bottom trawling on *Posidonia oceanica* meadows. *International Workshop on Posidonia oceanica beds*. Boudouresque C.F., Jedy de Grissac A. & Olivier, J. (edit.) *G.I.S. Posidonie* publ. 1 Fr: 63-72.
- PÈRES, J.M. 1984. La regression des herbiers a *Posidonia oceanica*. *International Workshop on Posidonia oceanica beds*. Boudouresque, C.F.; Jedy de Grissac, A. & Olivier, J. (edit.) *G.I.S. Posidonie* publ. 1 Fr: 445-454.
- RAMOS A.A. 1984. Cartografía de la pradera superficial de *Posidonia oceanica* en la bahía de Alicante (SE, España). *International Workshop on Posidonia oceanica beds*. Boudouresque, Ch. F.; Jedy de Grissac, A. & Olivier, J. (edit.) *G.I.S. Posidonie* publ. 1 Fr: 57-61.
- RAMOS *et al.* (in press).