

Expert System for Phytoplankton Classification (ESPHYNKS)

N. MARBA*, E. MERINO**, A. CRUZADO*, Z. VELASQUEZ* and J. PLUS**

*Centre d'Estudis Avançats de Blanes, C.S.I.C., Cami de Santa Barbara s/n, Blanes (Spain)
 **Facultat de Ciències Exactes, Ingeniería y Agrimensura, U.N.R., Av. Pellegrini 250, Rosario (Argentina)

The goal of the work reported here was to build a Prototype Expert System for the identification of marine phytoplankton.

The system runs PC-AT compatible computer, an auxiliary video monitoring, a TV camera and a digital video processing card. A commercially available inference tool has been used to facilitate the development task.

At the present stage of prototype development, ESPHYNKS allows identification of 8 phytoplankton genus and 16 species belonging to the genus *Coscinodiscus*. The knowledge base of the system is developed from a matrix (Fig.1) organized with a data base program. The *Coscinodiscus* matrix is composed of seven variables visible under optical inverted microscope, sufficient for unambiguous identification of each species. The data base program is used to explore the combinations of morphological characters for all *Coscinodiscus* species. After testing the matrix, combinations of characters are translated into expert system rules.

SPECIES	AREOLAE ARRANG	CENTRAL AREA	VALVES FORM (AV)	CONCAVE CENTRE	CHROMATOPHORES	THIN ABDELATION
C. CENTRALIS	RADIAL RINGS	WITH ROSETTE	CONVEX			NO
C. CONCENTRUS	RADIAL RINGS	WITH ROSETTE	CONVEX		IN ALL CYTOSOL	YES
C. CONCLUSUS	RADIAL RINGS	FACE AREA	CONVEX			
C. CURVATUS	GROUPED IN SECTIONS		CONVEX			
C. ECHINATUS	CURVED FANGENTIAL					
C. GRANIS	RADIAL RINGS	WITH ROSETTE	FLAT	YES		YES
C. GIGAS	RADIAL RINGS	FACE AREA	FLAT	YES		NO
C. LIMBUS	STRAIGHT FANGENTIAL					
C. MARGINATUS	PERIPHERAL AREA					
C. MODULIFER	RADIAL RINGS					
C. RETIATUS	FACE AREA	CAPILLAR				
C. OCCULUS IRIDIUM	RADIAL RINGS	WITH ROSETTE	FLAT	YES		NO
C. PERFORATUS	RADIAL RINGS	WITH ROSETTE	FLAT	NO		
C. PERFORATUS	RADIAL RINGS	FACE AREA	FLAT	NO		
C. RADIATUS	RADIAL RINGS	FACE AREA	FLAT			
C. STELLATUS	RADIAL RINGS	STAR LIKE FORM				
C. TUBIS	RADIAL RINGS	WITH ROSETTE	CONVEX		IN CENTRAL & MARGINAL REGIONS	YES
C. VALLISII	RADIAL RINGS	FACE AREA	FLAT	YES		YES

Fig.1 - Matrix organized with a data base program useful to build the ESPHYNKS knowledge base.

An image bank build from observations of field and laboratory phytoplankton samples under optical and scanning electron microscopy is associated to the knowledge base. During consulting, the system combines image display with questions to facilitate the identification. At the present time the image bank includes more than 450 different views of about 150 species.




Fig.2 - Reproduction of one of the screens on the auxiliary video monitor. This on presents eight phytoplankton genus included in the system.

ESPHYNKS structure provides different ways for species identification in field samples being observed under the microscope or described by the user. Each session begins presenting a screen (Fig.2) on the auxiliary video monitor with images of the eight phytoplankton genus included in the system. Two possibilities are considered depending on whether the sample is identical or not to one of the images. In the first case, the answer is trivial and the system concludes about the genus directly. In the second, an interactive process is established with the user by means of questions, help menus and images displayed on the video monitor leading to the complete determination of the genus. After this first phase is completed satisfactorily, the same procedure is repeated for that particular genus until the species is identified or not, depending on the answers given by the user. Further development of ESPHYNKS would consist on the repetition of the method applied in the construction of the prototype.

This novel computerized system proposes a new way for species classification and greatly facilitates the task to non specialists in the field of phytoplankton (e.g., technicians working in environmental assessment projects, students, etc.). The system is also an excellent tool for educational purposes. The use of relatively inexpensive hardware also allows for remote use of the system once it is fully developed. Experts and technicians around the world may send their own optical or SEM images on diskette or through electronic mail to a central identification centre where full identification could be accomplished.

Mobility of some Benzene Derivatives on Progressively Dried Fe (III)-Impregnated Silica Gel

Sonja ISKRIĆ*, Zeljka TOPOLČIĆ*, Nevenka ZAMBELI** and Olga HADŽIJA*

*Ruđer Bosković Institute, Zagreb, Croatia (Yugoslavia)

**Faculty of Pharmacy and Biochemistry, Zagreb, Croatia (Yugoslavia)

It is known that the surface of rocks, sediments and soils have active functional groups, in particular OH, derived from hydrous oxides or organic materials. The hydroxy groups of metal hydroxy/oxides are active factors for processes of interaction with organic ligands i.e. formation of inner sphere surface complexes (1). In the transformation of iron in soils and sediments, and also in biochemical systems, great importance has been attributed to the dissolution of iron hydrous oxides (2). In natural systems this process is affected by biogenic ligands such as organic acids - dicarboxylic or hydroxy carboxylic. Iron hydrous oxides may differ widely with regard to chemical composition and structure and consequently may have different chemical reactivity (3,4,5,6). In our work we examined the dependence of the level of hydration of iron hydrous oxides on the behaviour (binding) of some benzene derivatives, as a model system for natural conditions. As the technique thin-layer chromatography on silica gel impregnated with Fe(III)-ions was applied. The plates were prepared by spraying with iron nitrate solution and dried at 140°C for different time intervals. Developers were distilled and tap waters. Organic model compounds were pyrogallol and salicylic acid.

It was found that prolonged drying of the support increased mobility of the compounds. This phenomenon can be explained by formation of different species of silica gel Fe(III) complexes containing decreasing amounts at hydroxy groups and water. The results are given in Table 1. representing the dependence of $R_f \times 100$ of the compounds on drying time.

Table 1.

Compound	Time of drying at 140°C		
	5 min	3 h	22 h
Pyrogallol	$R_f = 20$	$R_f = 24$	$R_f = 54$
Salicylic acid	$R_f = 30$	$R_f = 37$	$R_f = 49$

References:

- W. Stumm, B. Wehrli and E. Wieland, *Croat. Chem Acta* **60** (1987) 429.
- W. Stumm, B. Sulzberger, Siffert, D. Suber and S. Banwart, *Mar. Chem.*, in press
- W. Schneider and B.B. Schwyn in *Aquatic Surface Chemistry*, W. Stumm Ed., Wiley and Sons, 1987, p. 167.
- S. Musić, A. Vertes, G.W. Simmons, I. Nagy-Czako and H.J. Leidheiser, *J. Colloid Interface Sci.* **85** (1982) 256.
- O. Hadžija, S. Iskrić and M. Tonković, *J. Chromatogr.* **402** (1987) 358.
- O. Hadžija, S. Iskrić and M. Tonković, *J. Chromatogr.* **464** (1988) 220.