

On the formation of the intermediate water masses off the Egyptian Mediterranean Coast

M.-A. SAID and A.-M. KARAM

National Institute of Oceanography & Fisheries, Kayet Bay, Alexandria (Egypt)

The presence of an intermediate water characterized by a secondary maximum of salinity in the Mediterranean Sea has a very peculiar phenomenon. Regions of formation of this water mass have been already more or less identified in the Mediterranean Sea. Along the Egyptian Mediterranean coast, the formation of this intermediate water mass was studied by Morcos (1972), who suggested that the area to the east of longitude 29° E and west of Alexandria is a secondary source of formation of the intermediate water. In addition, Abdel-Moati and Said (1987) suggested that the area in front of Damietta is another region of formation. The aim of the present work is to identify more accurately the locations of the formation of this water mass on the Egyptian Mediterranean shelf and its spreading using the isopycnic analysis.

The oceanographic data used were selected from several expeditions carried out by Egypt and different countries for the last 27 years (1959-1986). Water temperature and salinity data have been taken from 145 hydrographic stations in winter. The average values of temperature and salinity of these data were computed, using the optimum interpolation of the correlation algorithm, for stations distributed in a regular grid half degree latitude by half degree longitude for the winter season.

Our study confirms the findings of Morcos (1972) and Abdel-Moati and Said (1987). Besides, a new source has been identified which appears as a region of salinity > 39.30 ‰ (Fig.1) and a relatively high temperature > 17.00°C (Fig.2). The salinity distribution on the $\sigma_t = 28.80$ (Fig.1) indicates that the saline warm water flows away from the continental shelf to the northwest in a high-salinity tongue and it reaches the open sea with values as low as 38.95 ‰. This supports the suggestion of Morcos (1972) that more than one starting point representing various sources of formation of the intermediate water mass of maximum salinity are present in the Levantine basin.

The irregularities which appear particularly for the lateral salinity distribution west of 27° E (Fig.1) might be attributed to meso-scale noise in the data. In fact it needs more stress in the future work in order to get more clear picture of the thermohaline flow paths.




Fig.1. Salinity (‰) on the $\sigma_t = 28.80$ surface in winter.




Fig.2. Potential temperature on the $\sigma_t = 28.80$ surface in winter.

References

Abdel-Moati, A.R. and M.A. Said. 1987. Hydrographic structure of the Mediterranean shelf waters off the Egyptian coast during 1983-1986. *J. Thalassographica*, 10(2) : 23-39.
 Morcos, S.A. 1972. Sources of Mediterranean intermediate water in the Levantine Sea. In : *Studies in Physical Oceanography*, tribute to George Wüst on his 80th birthday, 185-206.

Contribution of the Rhone River water discharges to the dynamics of the Gulf of Lions in Autumn 1986

Jordi SALAT*, Antonio CRUZADO** and Michel PANOUSE***

*Institut de Ciències del Mar, Pasos Nacional s/n. 08003 Barcelona (Spain)

**Centre d'Estudis Avançats, Camí de Santa Barbara, 17300 Blanes (Spain)

***Laboratoire Arago, 66650 Banyuls de la Merendia (France)

The Gulf of Lions (NW Mediterranean) was covered by two oceanographic surveys, Pelagolion I and II, at the end of summer and at the end of autumn 1986. The main objective was to understand the role of the Rhone river water discharges on the pelagic production in that region at different seasons. Every cruise consisted on a series of CTD+Rossette casts scattered in the Gulf with a very fine survey in the vicinity of the river mouth. In this presentation, the results derived from CTD casts are summarized and discussed.

The main feature observed in the geostrophic velocity fields, obtained in both cruises (fig. 1), is the presence of the Liguro-Provençal-Catalan current contouring the shelf break, from east to west, with a maximum speed of about 35 cm/s, 15 miles south of Toulon (6°E) at the entrance to the region. This main current is present at all levels. Part of the water entering is deflected towards the river mouth along the north coast and another part flows to the south, mainly in summer, forming a cyclonic eddy. The main flow is minimum in the central part of the Gulf and it is enhanced again at the exit, directed towards the south 15 miles east of Cap Creus (42°20'N).

The main result concerning the dynamics is an anticyclonic eddy found in both cruises, in the northwest quadrant of the Gulf, centered on (43°N 3°30'E). River discharges produce a thin low salinity surface layer spreading over the vicinity of the river mouth. This water is trapped by the eddy and reconducted towards the coast allowing a long residence time and mixing. By this process, the initial thin surface layer of low salinity near the river mouth is converted to an homogeneous body of diluted water exposed to the local weather events in the coastal zone. This area acts as a reserve of the characteristic water type of the Gulf, one of the most important waters of continental influence present in the Western Mediterranean. For example, this reserve of diluted water, after strong cooling in central winter, and spreading through the bottom of the shelf break, as described by Fieux (1974) may be one of the sources of the subsurface salinity minimum found along the Catalan coast (Salat & Font, 1987).

At the beginning of summer, Allain (1960) found traces of such structures allowing the recirculation of the diluted water towards the coast. Therefore, by that season, the river runoff is usually much higher and the mixing layer above the thermocline is thinner than in autumn which allow a major spreading of surface low salinity layer as found by Castellón et al. (1985) in late spring. The summer observations of Millot (1981) show also an anticyclonic circulation in this area with local coastal upwelling events and downwelling in the center.

On the offshore side of the main current appears another anticyclonic eddy centered on (42°10'N 4°20'E) which seems to be very persistent. This eddy is completely uncoupled with the rest and the water in this region is not affected at all by the continental influence. Probably this eddy will play a major role on the deep water formation process during winter.




Figure 1. Geostrophic current and salinity fields at 10 m depth. a) at the end of summer 1986 b) at the end of autumn 1986

References

Allain C., 1960. Topographie dynamique et courants généraux dans le bassin occidental de la Méditerranée. *Rev. Trav. Inst. Sci. Tech. Pêches Marit.*, 24,1: 121-145.
 Castellón A., J. Salat & M. Masó, 1985. Some observations on Rhone fresh water plume in the catalan coast. *Rapp. Comm. int. Mer Médit.*, 29,3: 119-120.
 Fieux M., 1974. Formation d'eau dense sur le plateau continental du Golfe du Lion. *Colloques Internationaux du C.N.R.S.*, 215: 165-189.
 Millot C., 1981. La dynamique marine sur le plateau continental du Golfe du Lion en été. Thèse Doctorat. Univ Paris VI.
 Salat J. & J. Font, 1987. Water mass structure, near and offshore the Catalan coast during winter in 1982 and 1983. *Annales Geophysicae*, 5B,1: 49-54.