

O-VIII9

Double diffusive activities in the Cretan Sea during late Summer 1987

G. ZODIATIS

National Centre for Marine Research, Aghios Kosmas, 166 04, Athens (Greece)

The profiles of the hydrological characteristics (T,S) finestructure are examined in the Cretan Sea during the POEM-5 cruise, September - October 1987. CTD (SBE-9) data from 20 stations (Fig.1) averaged to 1 data per second of the original data which sampled at 33 Hz with a lowering speed of about 0.75 m/sec. The data interpolated with 1 dbar spacing and then are used for different finestructure analyses.




Fig.1 The study area showing the location of the hydrographic stations during the POEM-5 cruise.

One of the mechanisms that can generate the finestructure of the T-S characteristics is the double diffusive convection (Fedorov 1978, Karlin 1988 et al.). Generally is considered that the double diffusion and especially the salt fingering is an important process for the vertical transportation of heat and salt in the water column resulted by the different values of α and β coefficients (Turner 1985).

To find the nature and the strength of double diffusive activities, in the Cretan Sea, the parameter of Turner angle (Tu) is computed at depths between 100-500 dbars.

The observed profiles structure, at the majority of the examined depth interval, indicate that salt fingering process might be occurred. The hydrological conditions of the water at the above water columns are warm and saline water overlying colder less saline water, characterized as LIW of Cretan Sea. Turner angle profiles verify the prevalence of the salt fingering regime.

The stability regimes of double - diffusion (salt fingering, diffusive, stable, unstable), for all stations in the investigated area, are illustrated in the Turner angle (Tu) histogram (Fig.2). The stippled areas indicate the salt fingering and diffusive layering portions. The maximum Tu volume of 58% corresponding to the salt fingering while only 11% for the diffusive convection.

The above vertical distribution of Tu shows the predominance role of salt fingering process in the formation of the finestructure of Cretan water at intermediate depths.




Fig.2 Turner angle histogram with the stability regimes of double - diffusion for all stations in the Cretan Sea, late summer 1987.

REFERENCES

Fedorov K.N., 1978. The Thermohaline Finestructure of the Ocean, Pergamon Press Marine Series vol.2.
 Karlin L.N., Kloukov E.Y., Koutko V.P., 1988. Finestructure of the Hydrophysical Characteristics at the Upper Layer of Ocean, Hydrometeoizdat.
 Turner J.S., Convection in Multicomponent Systems, Naturwissenschaften, 72, N.2, 70-75pp.

O-IX1

Atmospheric Long Range Transport of pollutants to the North Western Mediterranean

Marta ALARCON*, Antonio CRUZADO* and Sergio ALONSO**

*Centre d'Estudis Avançats de Blanes, Blanes (Spain)

**Department of Physics, Group of Geophysical Fluid Dynamics, Balears Islands University (Spain)

Large quantities of substances of natural or anthropogenic origin enter the marine environment through the atmosphere, from both coastal and distant sources. This is especially true for semienclosed seas such as the Mediterranean Sea. Preliminary estimations of the fluxes of substances such as Hg, Cd, Pb and Cr, confirm that the atmospheric transport for these substances is, at least, comparable in magnitude to the riverine inputs (MED POL, 1989).

In a recent study (Alarcon and Cruzado, 1990) the nutrient fluxes (nitrate, nitrite, ammonia, phosphate and silicate) indicate that the deposition amounts are of the same order of magnitude to that of the vertical diffusive fluxes from deep waters into the photic zone. Therefore the atmosphere could be an important source of nutrients for the marine productivity, mainly in oligotrophic waters, as is the greatest part of the Mediterranean Sea.

The meteorology of the Mediterranean region is particularly variable both spatially and temporally, and this variability is due, partly, to the hilly orography. The topography behind the coastline around all the Mediterranean is complex. This can interact with the long range atmospheric flux inside the boundary layer influencing the horizontal transport and the vertical mixing of the substances by complex mechanisms.

At first, the knowledge of the deposition processes of the aerosol particles, combined with transport models should allow an evaluation of the net transfer rate from air to sea of particulate matter.

Computation of air parcel trajectories is a very powerful tool to estimate the long-range transport of substances. In this study we have computerized an isentropic method due to Petersen and Uccellini (1979) to simulate long-distance transport of substances.

The isentropic approximation reduces a tridimensional trajectory computation to a bidimensional problem. The air parcel moves above isentropic surfaces. Although only the horizontal components of the wind are used, the vertical velocity of the particle is not ignored. It is implicitly taken into account through the contour and the temporal variation of the isentropic surface.

Trajectories are computed applying an explicit system of equation based in the theory of the "discrete model" developed by Greenspan (1972), to the atmospheric equations of the movement. Trajectories are constructed in an isentropic framework, from the movement equations, where the pressure fields are represented by the Montgomery stream function, $Cp+gz$.

The model calculates parcel positions in time steps of 900 s. A trajectory segment is integrated by 48 time steps. Acceleration is computed from the gradient of the Montgomery streamfunction, and then velocity and position are interpolated for the next time step. The process is repeated for the successive time steps to complete a 12 hours trajectory segment.

The stability of this process is discussed by Petersen and Uccellini (1979), and conclude that this explicit method is stable for time increments between 300 and 1800 s.

REFERENCES

Alarcon, M and Cruzado, A; (1990): Nutrient inputs into the North-Western Mediterranean Sea. EROS 2000, second workshop, Blanes.
 Greenspan, D; (1972): A new discrete mechanics with applications. J. Franklin Ins., 294, 231-240.
 MED POL; (1989): Airborne pollution of the Mediterranean Sea. MAP Technical Reports Series No.31.
 Petersen, R and Uccellini, L W; (1979): The computation of isentropic atmospheric trajectories using a 'discrete model' formulation. Mon. Weather Rev., 107, 566-574.