

## IAEA - 307 : A New Biological Reference Material for Marine Environmental Radioactivity Studies

A. VEGLIA\*, S. BALLESTRA\*\* and D. VAS\*\*

\*Centre Scientifique de Monaco, 16 boulevard de Suisse (Monaco)

\*\*International Atomic Energy Agency, International Laboratory of Marine Radioactivity, 19 avenue des Castellans (Monaco)

A laboratory intercomparison exercise to determine the levels of natural and artificial radionuclides in a homogenized sample of sea plant (*Posidonia oceanica*) was organized by the International Laboratory of Marine Radioactivity (IAEA) in 1988-1989. The sea plants were collected along the shore from the vicinity of Monaco in October 1986 and were assumed to be contaminated by both global and close-in radioactive fallout from the accident of Chernobyl (26 April 1986).

TABLE 1

## Reference activity values :

Radionuclide	Recommended value <sup>1</sup> Bq kg <sup>-1</sup>	Confidence interval Bq kg <sup>-1</sup> ( $\alpha=0.05$ )
Potassium-40	150	141-161
Ruthenium-106	33.5	30.0-36.5
Silver-110m	5.1	4.8-5.5
Caesium-134	1.6	1.5-1.9
Caesium-137	4.9	4.5-5.2
Radium-226	3.1	2.1-4.4
Plutonium-238	0.025	0.022-0.028
Plutonium-239,240	0.72	0.66-0.79
Americium-241 <sup>2</sup>	0.036	0.030-0.050

1. Activities are expressed on dry-weight basis (constant weight at 80°C) for the reference date : 1 January 1988.

2. Recommended value for reference date only. Ingrowth from any <sup>241</sup>Pu present will alter value if <sup>241</sup>Am is measured in the sample during some subsequent year.

Sixty-six laboratories from 31 member state countries reported results for 34 natural and man-made radionuclides in this reference material (IAEA-307). The large participation made it possible to generate statistically valid recommended values for the activities of 9 radionuclides in this material (Table 1). Non-certified information values were generated for 4 radionuclides (Table 2). This reference material IAEA-307 was certified as a reference material from this exercise and made available for distribution in 1990.

TABLE 2

## Non-certified activity values :

Radionuclide	Activity <sup>1</sup> Bq kg <sup>-1</sup>	Confidence interval Bq kg <sup>-1</sup> ( $\alpha=0.05$ )
Strontium-90	0.72	0.28-1.6
Lead-210	58.5	40-91
Thorium-228	3.2	1.5-4.2
Uranium-238	14	2.5-21

1. Activities are expressed on dry-weight basis for the reference date : 1 January 1988.

Recommended concentrations represent overall median values calculated from the results that passed a non-parametric test that identified outliers (1) and which satisfied a number of other selected quality criteria (2). Confidence intervals were calculated from statistical tables and correspond to a significance level of 0.05.

The total number of outliers from the 415 reported results was small (about 8% of all results) and varied between 0 and 3 per laboratory. Twenty laboratories submitted results that included at least one outlying value.

The reference material IAEA-307 is certified for use to evaluate the accuracy of radioanalytical procedures for biological materials, and when checking precision among analysts in a laboratory.

## REFERENCES

- 1) A. Veglia : International Atomic Energy Agency, Report No. IAEA/RL/84 (August 1981)
- 2) A. Veglia, S. Ballestra, D. Vas : Report on the Intercomparison of IAEA-307, Radionuclides in Sea Plant, Report No. IAEA/AL/014 (October 1989)

## Gamma Emitters in IAEA's SD-A-1 Marine Sediment Intercomparison Sample

Iolanda OSVATH



Environmental Radioactivity Research Laboratory, Institute of Meteorology and Hydrology, Bucharest (Romania)

SD-A-1 is a deep-sea sediment sample, collected in 1985 from the Atlantic floor (between 46° - 48°30' N and 18° - 22° E, depth greater than 4000 m), (IAEA, 1988). A box corer was used for sampling. The top 15 cm of the resulting sample having been removed, the remaining material was assumed to have artificial radionuclide concentrations at or below detection limits. The intercomparison run, organized by the International Laboratory of Marine Radioactivity during 1986-1987, certified the values of some natural radionuclide concentrations in the sample, which thus became reference material, also recommended as a blank for artificial radionuclides. Results for K-40 and Ra-226 are synthesized in Fig. 1, respectively Fig. 2.

In our laboratory, we have analysed an aliquot of 100g of the sample, using high resolution, low-background gamma spectrometrical equipment. Three measurements were performed, with the counting time ranging between 30 and 61 hours. Data were processed according to ILMR requests.

Two problems regarding data interpretation ask for attention, leading to conclusions of interest for the gamma spectrometrical analysis of sediment samples.

1. Among the Ra-226 concentration values reported by the participants, the ones obtained through gamma spectrometrical analyses were assessed by two methods:  
a. directly, from the 186 keV photopeak, and  
b. indirectly, from Pb-214 - Bi-214, assuming them to be at equilibrium with Ra-226. Under normal sealing conditions (sample in closed plexiglass beaker), even at times much longer than needed for equilibrium to be attained among Ra-226 daughters, a disequilibrium factor of 1.3-4 persists. It is due to the great diffusivity of Ra-222,

Fig. 1. Reported values for KALIUM-40 : mean  $\pm$  standard deviationFig. 2. Reported values for RADIIUM-226 : mean  $\pm$  standard deviation

part of which can consequently leave the sample. This is not the case with the thorium series, due to the relatively short half-life of Ra-220, the Ra-228(Ac-228)/Pb-212 activity ratio being very close to unity. These statements are based on a large amount of measurement data resulting from marine as well as river and lake sediments.

The conclusion can be drawn that the activity concentration of the Ra-226 in the sample cannot be assimilated to the Pb-214 - Bi-214 activity concentration.

2. Activity concentrations are reported for Tl-208, which appear to be calculated using different values for the yield of its gamma line at 583.1 keV. According to the disintegration scheme of the thorium series (GUSEV and DMITRIEV, 1978), when equilibrium is reached in a closed sample, as pointed out in pp. 1, the Pb-212 - Bi-212/Tl-208 activity ratio approximates the branching ratio (BR=0.36).

This BR has to be considered when calculating the Tl-208 activity concentration from its 583.1 keV line.

The same explanation also accounts for the large scatter in Tl-208 data reported in the IAEA-306 Baltic Sea sediment intercomparison run (IAEA, 1989): they obviously group around two values, which differ from one another by a factor which closely approximates the BR.

Regarding U-235, although itself was one order of magnitude below our detection limit, the sample spectra showed clearly the presence of its gamma emitting daughters, like Th-227, Ra-223 and Bi-211. Most of the important gamma lines of the daughters are usually shielded by those of radionuclides in the U-238 and Th-232 series.

## REFERENCES

- IAEA, 1988. Report on the Intercomparison Run SD-A-1: Radionuclides in Marine Sediment. IAEA/AL/012  
IAEA, 1988. Report on the Intercomparison Run IAEA - 306: Radionuclides in Baltic Sea Sediment. IAEA/AL/013  
GUSEV I.G., DMITRIEV P.P., 1978. Radioactive Series. A guide.