

# O-IX8

## Computation of the Surface Heat Flux by means of bulk formulas in El-Max Bay, Alexandria (Egypt)

M.A. SAID

National Institute of Oceanography & Fisheries, Kayet Bay, Alexandria (Egypt)

El-Mex Bay, west of Alexandria, extends for about 15 km and has a mean depth of 10m. Its surface area is of about 19.4 km<sup>2</sup> and its volume 190.3 x 10<sup>6</sup> m<sup>3</sup>. It receives a heavy load of wastewater (2.4 x 10<sup>9</sup> m<sup>3</sup> / year) both directly from industrial outfalls or indirectly from Lake Maryut via El-Mex Pumping Station.

Throughout the period from January to December 1988, 12 trips were carried out in El-Mex Bay area using a motor boat. Temperature and salinity were measured at discrete depths from seven hydrographic stations. In particular, monthly data were there given as function of depth describing the mass field characteristics of the water column through a year of 1988. The monthly mean climatological data from Ras El-Tin Meteorological Station at Alexandria for the 15 years (1970-1984) have been used for the computations of the heat flux.

The total upward heat flux  $Q$  from the sea surface consists of the upward flux of long wave radiation  $Q_L$ , latent heat flux  $Q_e$  and the sensible heat flux  $Q_c$  less the absorbed global irradiation  $Q_s$  :

$$Q = Q_L + Q_e + Q_c - Q_s \quad (1)$$

Formulas for computation of  $Q_L$ ,  $Q_e$  and  $Q_c$  are taken from Stravisi and Crisciani (1986) which are given according to Gill (1982). A detailed discussion of the formulas is presented in Stravisi and Crisciani (1986). Rate of absorption of global solar irradiation  $Q_s$  is given by

$$Q_s = (1 - A_n) Q_n$$

where  $A_n$  is the sea surface albedo and can be calculated from tables (Kolesnikov, 1970).  $Q_n$  is the global solar irradiance calculated at the sea level using Timofeev's equations (1970). The heat flux (1) released from the sea surface is the difference  $Q = Q_b - Q_\theta$  between an input flux  $Q_b$  through the other boundaries, mainly due to advection,

$$\text{and} \quad Q_\theta = C_w \int h \frac{\partial \theta}{\partial t}$$

representing the rate of heat storage in the sea, in a vertical water column of depth  $h$  and unit horizontal surface;  $\theta$  is the vertical mean seawater temperature,  $C_w = 3.98 \times 10^3 \text{ JKG}^{-1} \text{ K}^{-1}$  is the average seawater specific heat at constant pressure. The boundary heat flux  $Q_b$  in a basin can be computed once  $Q$  and  $Q_\theta$  have been evaluated.

The monthly mean values of the computed heat flux are listed in table 1.

Table (1). Monthly and annual surface heat fluxes (Watts / m<sup>2</sup>) from El-Mex Bay.

	January	February	March	April	May	June	July	August	September	October	November	December	Year
$Q_L$	76.13	79.55	82.97	83.34	79.18	72.86	66.24	64.42	66.95	69.12	71.64	73.79	73.85
$Q_e$	96.05	100.58	108.58	109.48	115.91	130.42	148.67	152.17	149.93	119.43	103.45	103.27	119.83
$Q_c$	18.20	10.90	-0.84	-7.70	-9.46	-9.03	-7.28	-4.40	00.00	3.55	7.91	16.93	1.58
$Q_s$	109.81	138.21	223.82	290.67	315.48	368.29	355.87	318.55	255.03	186.90	132.28	103.56	234.87
$A_n$	0.080	0.073	0.068	0.061	0.062	0.054	0.054	0.054	0.060	0.068	0.077	0.082	0.066
$Q_s$	101.02	146.66	209.05	272.94	295.92	348.40	336.65	303.35	239.73	174.19	122.09	95.07	220.26
$Q$	89.38	44.37	-18.14	-87.82	-110.29	-154.15	-129.02	-89.16	-22.85	17.91	60.91	98.92	-25.00
$Q_\theta$	-18.00	24.98	15.64	24.97	61.66	56.10	21.20	-15.36	-28.40	-21.40	-52.11	-59.28	00.00
$Q_b$	71.38	59.35	-2.50	-62.85	-48.63	-98.05	-107.82	-104.52	-51.25	-3.49	8.80	39.64	-25.00

For a better comprehension of the "heat sink" character of El-Mex Bay, the flux  $Q_\theta$  of heat stored in the Bay has been computed, using a mean depth  $h = 10\text{m}$ , together with the heat  $Q_b$  advected from the Bay (Fig.1). The heat storage values vary from -59.28 in December to 61.66 W m<sup>-2</sup> in May. The flux  $Q_b$  is negative during the period from March to October with an average of -59.89 W m<sup>-2</sup>. From November to February, it varies between 8.80 and 71.38 W m<sup>-2</sup>.




Fig.1. Annual cycle of the heat storage  $Q_\theta$  and the advected heat flux  $Q_b$  in El-Mex Bay.

### References

- Hill, A.E.1982. Atmospheric-Ocean dynamics. Acad. Press, New York,662p
- Kolesnikov, A.Q.(ed).1970. Atlas of the heat balance of Oceans. Morsk. Hidrofiz. Inst. : Sevastopol. (In Russian).
- Stravisi, F.& F. Crisciani.1986. Estimation of surface heat and Buoyancy fluxes in the Gulf of Trieste by means of bulk formulas. Bollett. di Oceanologia Teorica ed Applicata, 6 : 55-61.
- Timofeev, N.A.1970. On the external components of the heat balance in the oceans. Marine hydro.inves.,No 1, Sevastopol, 148-165 (In Russian)