

## B-III1

### Amphipods and Molluscs of the circalittoral enclaves onto dead terraces of degraded *Posidonia oceanica* Meadows on the Coast of Alboraya (Spain, Gulf of Valencia, Western Mediterranean)

I.-M. GINER, A. MARTI, V. BENEDITO, J.-L. ESTEBAN, J. TORRES, A.-M. GARCIA-CARRASCOSA and R. CAPACCIONI

Invertebrates and Marine Biology Laboratory, Department of Animal and Cellular Biology and Parasitology, Faculty of Biological Sciences, Valencia University, 46100 Burjassot, Valencia (Spain)

The epigraph of circalittoral enclaves refers to those structures originated by sciaphilic biogenous accretion onto soft substrate rhizomes or *P. oceanica* rhizomes remains.

On the coast of Alboraya, and in general in the Gulf of Valencia, the ascension of circalittoral biocoenosis towards shallower waters can be observed. This ascension is due to water turbidity levels, rising as a result of high anthropic pressure that this zone presents (urban, industrial, turistic and agricultural activities). In the studied area these concreted masses use dead terraces of degraded *P. oceanica* meadows for their instalment and the lower limit of their bathymetric range oscillates between -10 and -12 m.

The installation of these structures on *P. oceanica* meadows occurs, as is described by (18), in those deep meadows with medium leaf density or in shallow ones with high leaf density, where sciaphilic environments can be found. However the *P. oceanica* meadows, in Alboraya, presents a high regression degree with a very low shoot density (1-2 shoots/m<sup>2</sup> at -10 m) and is therefore unable to create sciaphilic biotopes on its own. Then the reason why these concreted masses appear at shallow levels is because of the turbidity conditions of the sea-water. In fact, in shallower areas (2-3 m depth) Secchi disk disappears at 1m depth and in even offshore deeper areas (around -17 m) it does at 6-7 m.

These structures rise 30-40 cm from the bottom, and they are more frequent on the corriches that delimit the pot-holes and channels. Concreted masses are built by the action of calcareous algae (*Pseudolithophyllum expansum*, *Lithophyllum mammosum* and *Mesophyllum lichenoides*) and by hard structures of porifera and bryozoans, which include shell remains, sediment of diverse texture. *P. oceanica* rhizoma fibril remains, etc.. They are covered by an important sciaphilic flora (*Peyssonnelia* sp., *Udotea petiolata*, *Halimeda tuna*, *Codium bursa*, *Sphaerococcus coronopifolius*, etc.) and fauna (*Eunicella cavolinii*, *Pentapora fascialis*, *Myriapora truncata*, *Halocynthia papillosa*, etc.).

Amphipod fauna, studied by (13), shows a first stock which is formed by the species *Iphimedia serratipes*, *Lysianassa pilicornis* and *Pseudoprotella phasma* that have been previously mentioned from coralligenous bottoms 9, 11, 19. The second stock is constituted by *Microdeutopus algicola* a species with affinity to hard substrates with vegetal coverage. The third one is represented by species with wide ecological distribution: *Gammarella fucicola* (1) (5) (6) (10) (11) (14), and *Corophium sextonae*, which is presented in the whole biocoenoses at the studied zone except in SGCF. *Leucothoe richiardii* and *Atylus massiliensis* form a stock of species that come from biotopes around. The former comes from rhizome terraces of *P. oceanica* (3) (8), and the latter from sandy biocoenoses (2). The last stock is constituted by a form of *Maera* sp. whose peculiar features don't allow us to assign it to any of the nine species known from the Mediterranean (7). In the studied area, *Maera* sp. appears exclusively with an important density on these enclaves.

Referring to molluscan fauna, described by (4), there is a first group formed by *Cianculus cruciatus*, *Diodora graeca*, *Raphitoma echinata*, *Turbona cimex*, *Columbella rustica*, *Chauvetia minima*, *Chama gryphoides* and *Muricopsis cristata* previously mentioned in the Coralligenous (18) (20) (15) (16). The second stock is formed by greater numerous species distributed fauna (*Tricolia pullus*, *Rissoa violacea*, *Jujubinus exasperatus*, *Cianculus jussieuvi*, *Venerupis pullastra*, *Glans trapezia*, etc.) characteristic of *P. oceanica* meadows. The third stock shows a faunistic group characterized by species with affinity to hard substrates (*Arca noae*, *Musculus costulatus*, *Gastrochaena dubia* and *Lithophaga lithophaga*).

The colonization of circalittoral populations in less deep zones is demonstrated by the localization of these shallow circalittoral biogenous enclaves, and by the unusual existence between -6 and -11.5 m at the studied area, of important populations of the amphipod *Lembos angularis* species characteristic of deeper muddy bottoms (8) (3).

In the present study the lower limit of *P. oceanica* meadows, located at -25 m fifteen years ago, have been found at 17-18 m depth. This ascension of the lower limit is probably due to the increment of the turbidity conditions mentioned above. These higher sciaphilic conditions and the existence of dead terraces of *P. oceanica* rhizomes, have conditioned the gradual rising of Coralligenous towards shallower depths. In fact, wide zones with coralligenous blocks of about 2 m height, forming continuous strings, are found up to the -20 m isobat.

#### BIBLIOGRAPHY.

- (1) BELLAN-SANTINI (D.), & LEDOYER (M.), 1973.- *Téthys*, 4 (4), pp.399-934.
- (2) BELLAN-SANTINI (D.), KARAMAN (G.), KRAPP-SCHICKEL (G.), LEDOYER (M.), MYERS (A. A.), RUFFO (S.) & SCHIECKE (U.), 1982.- *Mém. Inst. Oceanogr. Monaco*, 13, pp.1-364.
- (3) CHEVREUX (E.), 1910.- *Mém. Soc. Zool. Fr.* 23, pp.145-285.
- (4) GINER (I.M.), 1989.- Tesis de Licenciatura, Facultad de Biológicas, Universitat de Valencia. 225p.
- (5) HARMELIN (J.G.), 1964.- *Rec. Trav. Sta. mar. Endoume*, 51 (35), pp.43-106.
- (6) JACQUOTTE (R.), 1962.- *Rec. Trav. Sta. mar. Endoume*, 44 (29), pp.27-42.
- (7) KARAMAN (G.) & RUFFO (S.), 1971.- *Mem. Mus. civ. Stor. nat.*, Verona. 19, pp.113-176.
- (8) KRAPP-SCHICKEL (G.), 1975.- *Boll. Mus. civ. Stor. nat.*, Verona, 11, pp.91-118.
- (9) KRAPP-SCHICKEL (G.), 1976.- *Bull. Zool. Mus. Univ.*, Amsterdam, 5 (5), pp.31-45.
- (10) LEDOYER (M.), 1962.- *Rec. Trav. Sta. mar. Endoume*, 25 (39), pp.117-235.
- (11) LEDOYER (M.), 1968.- *Rec. Trav. Sta. mar. Endoume*, 60 (44), pp.125-295.
- (12) LEDOYER (M.), 1970.- *Bull. Inst. oceanogr. Monaco*, 69 (1406), pp.1-32.
- (13) MARTI (A.), 1989.- Tesis de Licenciatura, Facultad de Biológicas, Universitat de Valencia, 152p.
- (14) MASSE (H.), 1962.- *Rec. Trav. Sta. mar. Endoume*, 42 (27), pp.221-259.
- (15) PEREIRA, 1980.- *Com. Prim. Congr. Nac. Malac.*, Madrid, pp.79-84.
- (16) PEREIRA, 1985.- *Actas II Simp. Ibér. Estud. Bentos Mar*, 3, pp.243-251.
- (17) PÉRES (J.M.), 1967.- *Oceanogr. Mar. Biol. Ann. Rev.*, 5, pp.449-533.
- (18) PERES (J.M.) & PICARD (J.), 1964.- *Rec. Trav. Sta. Mar. d'Endoume*, 31 (47), pp.5-137.
- (19) RUFFO (S.) & SCHIECKE (U.), 1979.- *Boll. Mus. Stor. nat.*, Verona, 5, pp.401-429.
- (20) SPADA (G.), SABELLI (B.) & MORANDI (V.), 1973.- *Conchiglia*, 9 (3-4), pp.29-67.

B