


Late Pleistocene Paleoclimates and Anoxic Events in the Eastern Mediterranean : The Deep-Sea Record

Yvonne HERMAN

Department of Geology, Washington State University, Pullman, WA 99164 (U.S.A.)

Faunal composition and oxygen isotopic analyses of foraminiferal and pteropodal shells were utilized to reconstruct the paleoclimatic and paleohydrologic history of the Eastern Mediterranean. The reconstruction was done by mapping the recent fauna in core "tops" deposited during the Holocene and their calibration against observed present-day temperatures and salinities in the water column. The broad data base, using published faunal distributions from the world ocean, covers a wider range of temperatures and salinities and a combination of these factors, than those which are thought to have existed during glacial periods in the Mediterranean. In the longest core with a nearly continuous record, spanning ~ 500 K years is contained in the ~ 1000 cm long Lamont-Doherty Geological Observatory (LDGO) core RC9-181. Within this time interval represented by RC9-181 six major cold-warm cycles, correlatable to Emiliani's isotopic stages 1-12 occur. Calcareous



nannoplankton biostratigraphic and biochronologic framework was utilized for determining the times various events occurred and for estimating rates of sedimentation. Two important datum levels were recorded: the extinction of *Pseudoemiliana lacunosa* between 899 and 938 cms (0.44-0.46 Ma ago), and the first appearance of *Emiliana huxleyi* 0.26-0.27 Ma ago, between 455 and 479 cm depth in core. During glacial temperature minima, surface water temperatures were ~ 3°C lower in summer and 3-4°C lower in winter. Stadial and interstadial salinities were variable, reaching highest values during the last glacial temperature minimum when climates were more arid than today, sea level stood very low, the Nile discharge was greatly reduced and the connection between the Mediterranean and the Black Sea, which is a major supplier of low salinity water was severed. Following global warming and subsequent massive deglaciation, sea level rose. When the sea stand reached the Bosphorus sill (~36 m) the connection between the Mediterranean and the Black Sea was reestablished and the low salinity Black Sea water spilled over into the Mediterranean. A significant increase in precipitation and river runoff is also recorded during transitional climatic periods. These compounded effects produced a low density surface water layer which restricted thermohaline convection. The result was stagnation of the subsurface water and subsequent deposition of sapropels, some of which were laid down during intervals of pronounced density stratification. Surface water salinities dropped to low values during the deposition of sapropels as evidenced by oxygen isotope data. Twelve sapropels, characterized by distinct faunal associations, documenting different degrees of stagnation, occur in sediments representing the last 0.5 Ma.

TABLE 1 Location, depth and length of cores

Core	Latitude (N)	Longitude (E)	Depth of water (m)	Length of core (cm)
V10-65	34°37.00'	23°25.00'	2586	960
V10-67	35°42.00'	20°43.00'	2890	830
RC9-181	33°25.00'	25°01.00'	2286	930
RC9-185	34°27.10'	20°07.00'	2858	902

ACKNOWLEDGEMENTS.

I thank Lamont-Doherty Geological Observatory (LDGO) for making the R. Conrad and Vema cores available, and the Core Curators, most recently Rusty Lotti, for sampling and shipping the core samples to WSU. Support for the curating facilities of the LDGO Deep-Sea Sample Repository was provided by the NSF through Grant OCE85-00232 and the ONR through Grant N00014-84-C-0132. This investigation was supported by the donors of the Petroleum Research Fund administered by the American Chemical Society Grant 10939-AC2.