

Characterization and Distribution of Organic Matter in recent sediments from Elefsis Bay (Saronikos Gulf, Greece)

Ulrich MANN* , Detlev MULLER* and Christos ANAGNOSTOU**

*Institute of Petroleum and Organic Geochemistry (ICH-5) at the Research Center (KFA) Juelich, P.O. Box 1913, D-5170 Juelich (F.R.G.)

**National Center for Marine Research (NCMR), Aghios Kosmas, 16604 Hellinikon (Greece)

Within Recent sediments from Elefsis Bay, two types of geogenic and two types of anthropogenic organic matter may be expected:

- 1: From autochthonous, biogenic marine organisms like phyto- and zooplankton as well as from benthos.
- 2: Wind- or river transported plant debris (spores, pollen, plant tissues).
- 3: Petroleum contamination from oil refineries, shipyards and harbour areas as well as other organic compounds from industrial sources.
- 4: Sewage effluents from the Athens metropolitan area.

The objective of our investigations is to provide an organic-geochemical survey of the sediments from Elefsis Bay, to identify the four above mentioned organic matter types and to establish the presence of potential anomalous organic concentrations.

Sediment samples were taken in March and October 1989 using a modified Reineck corer on board the Greek R/V "Aegeio". Recovered sediments constitute about 80% clay plus silt, in which approximately about one half is represented by carbonate and the other half predominantly by clay minerals.

Many of our basic organic-geochemical results confirm the findings by VOUTSINOU-TALIADOURI & FRAGOUDAKI (1986) who determined total organic carbon contents (TOC) of up to 4%.

Total content of soluble organic matter, as well as relative amounts of aliphatic to aromatic hydrocarbons to heterocompounds (N,S,O), do not exhibit any significant variation, most probably due to the various sources of organic matter within Elefsis Bay. Only the concentration of aromatic hydrocarbons seem to monitor the proximity of industrial effluents.

The individual types of organic matter were elucidated as follows:

Macro- and microscopic observations indicate that the autochthonous biogenic, marine constituents of the sediments from Elefsis Bay are nearly exclusively made up of carbonate shells. Because shell particles are concentrated with the sand fraction, their regional contribution to the sedimentary input can be estimated via the carbonate content. The amount of preserved organic matter, however, has to be identified via the total organic carbon content of the sand fraction. As we have observed, most pelecypods and gastropods preserve at least some of their organic matter within their shells.

Relative amounts of the marine versus terrestrial organic matter present within the sediments were evaluated by concentrations of the C₁₅ to C₁₈ versus C₂₁ to C₂₅ aliphatic hydrocarbon fractions. Within the sediments from Elefsis Bay, terrestrial plant material predominates. This is in accordance with the very close shoreline and the wind direction from the inland area. Nevertheless it can also be argued that anthropogenic contamination of Elefsis Bay has reduced the quality of living conditions for marine organisms and therewith the primary availability of marine organic matter. The stress factor within Elefsis Bay is especially expressed by the scarcity of marine species.

As outlined above, a major contamination in Recent Elefsis Bay sediments is represented by petroleum together with other organic compounds. Within all anoxic sediments we found an "Unresolved Carbon Mixture" (UCM) in the C₁₅₊-aliphatic hydrocarbon fraction which seems to correlate to an UCM within the C₁₅₊-aromatic hydrocarbon fraction. At the reference site outside Elefsis Bay, an UCM could still be identified, however, with a considerable lower concentration. No UCM was observed within the underlying oxic sediments.

Preliminary investigations reveal that the UCM is confined to the clay fraction of the sediment, especially to its adsorption capacity as expressed by the size of specific surface area.

The effects of sewage effluents within Elefsis Bay have not been evaluated so far. However, for this purpose, analyses of low-molecular-weight hydrocarbons are in progress.

The incorporation and preservation of marine, as well as of terrestrial organic matter, in Recent Elefsis Bay sediments, has been demonstrated from the results of our organic-geochemical survey.

Within anoxic sediments, organic, anthropogenic contamination is degraded to a considerably lower extent than within oxic sediments. As organic contamination seems to be strictly tied to clay minerals, we recommend detailed physico-chemical analyses to evaluate the chance and importance of potential remobilization processes.

References:

VOUTSINOU-TALIADOURI, F. and FRAGOUDAKI, St., 1986. Geochemical studies in sediments from a semi-enclosed polluted basin. Elefsis Systems Project Technical Report No.3, N.C.M.R., 12pp.