

Ichthyoplankton of the Egyptian Mediterranean waters IV- Distribution and occurrence of Mullet Larvae

N.-M. DOWIDAR and H.-H. EL-RASHIDY

Oceanography Department, Faculty of Science, Alexandria University, Alexandria (Egypt)

The seasonal and spatial distribution and abundance of mullet larvae and fry along the Egyptian Mediterranean coast were studied during the period from January 1982 to October 1984. The stations sampled and the study area are described by El-Rashidy and Dowidar (1990) in this volume.

A total of 859 mullet larvae occurred in the ichthyoplankton samples collected throughout the period of study. The larvae of *Mugil cephalus* and *Liza saliens* were observed in plankton samples throughout the period from July to November. As judged from the length frequency of the recorded larvae, it may be concluded that spawning of both species may begin during June and ends in late October - early November. The surface water temperature during this period varied between 21.5° and 28.4° C. These results are in accordance with the spawning seasons determined for both species by various authors from the study of gonad maturation (Rafail, 1968; Abdel Hamid, 1969; Youssef, 1973). Figure (1-A) describes the abundance and spatial distribution of the larvae of *M. cephalus* and *L. saliens* which revealed that they mostly spawn in coastal waters particularly in the eastern area (Burullus - Arish), at depths ranging from 20 to 50 m, and a distance of 3.5 - 10 km from the coast.

The larvae of *Liza ramada* were recorded throughout the period from November to April, contributing about 94% of the mullet larvae recorded during November. The length frequency of the larvae may indicate that the breeding of *L. ramada* begins in November and probably ends in March with the peak in late November to early December. The surface water temperature varied between 17.7 and 21.5° C. This agrees with the spawning time given by other authors working on the gonad maturation of the fish (El-Sedfy, 1971; Youssef, 1973; El Maghraby et al., 1974). The pattern of distribution of *L. ramada* larvae (Figure 1 :B & C) shows that the small larvae up till 7 mm were dominant in the offshore and middle zones. This indicates that *L. ramada* spawns at a distance of 15 - 27 km from the coast covering depths from about 50 to 200 m, during November and December. At the end of the spawning season in February, the larger fry (19 - 29 mm) were recorded in the coastal waters of Damietta and El-Diba, i.e. attracted to shallower depths at drain outlets and estuaries with relatively lower salinities. Our study reveals that *M. cephalus* larvae were less abundant than those of *L. ramada*; the later species constituted about 90% of all mullet larvae recorded. Such depletion of *M. cephalus* larvae may be attributed to the intensive fishing of these larvae for raising in fish farms. Approximately 20 million fry are collected annually from the coastal waters adjoining fresh and brackish water outlets particularly from El-Mex area. This process, in addition to fishing of the sexually mature fish during their spawning migration from the delta lakes to the sea, has undoubtedly exhausted the mullet stock in the Egyptian Mediterranean waters. Further more the increasing rate of pollution of the coastal waters particularly in the areas of larval attraction may affect the larval occurrence.




Figure 1 : Distribution and abundance of the different size groups of total mullet larvae.

References :

Abdel Hamid, Kh. 1969. M.Sc. Thesis, Faculty of Science Alexandria Univ.
 El-Maghraby, A.M., M.T. Hashem And H.M.-El Sedfy, 1974. Bull. Inst. Ocean. & Fish., A.R.E. 4 : 3 - 31.
 El-Sedfy, H.M., 1971. M.Sc. Thesis, Faculty of Science, Alexandria Univ.
 Rafail, S.Z., 1968. Stud. Rev. Gen. Fish. Coun. Mredit. 35:1-19.
 Youssef, F.S., 1973. M.Sc. Thesis, Faculty of Science, Cairo Univ.

The Development Rates of European Pilchard (*Sardina pilchardus* Walb. 1792) Eggs

Bülent CİHANGİR

Dokuz Eylül University, Institute of Marine Sciences and Technology, P.O.Box 478. 35213 Izmir (Turkey)

Using a Bongo-Net, pilchards eggs were collected at the peak of the spawning in Izmir Bay in February 1990. The eggs were roughly separated from plankton on board and transported to land laboratory as soon as possible in 6°C. The eggs were sorted according to their stages under dissecting microscope in the laboratory. The earliest stages were Ib 2 taken to three different temperature regimes (13 - 16 - 19 °C). It was estimated that the spawning time of pilchard is between 1900 and 2100 hours (PEREZ and RODRIGUEZ, 1988 ; CİHANGİR (in prep.)). Eggs have been sampled during twilight (1600 - 1900 h.), the sea water temperature was 14 °C at 20 meter depth. It was assumed that youngest captured eggs were 15-18 hours old. In laboratory experiments, hatching occurred in 75 h at 13°C, in 60 h at 16°C and in 55 h at 19°C (Fig.1) (the incubators were fluctuated to 0.5 - 1.0°C during the experiment). These results are approximate to RUSSEL (1976).

Stages description were adapted from MOSER and AHLSTROM (1985) and ALHEIT et al. (1987).





Figure 1. Development rates of pilchard eggs under different temperature regimes.



Various stages of pilchard eggs.

Description of Stages

- IIa 1 From fertilization to 64-cell stages.
- IIa 2 Formation of the blastodisc.
- IIa 3 Formation of the blastodisc as a lens.
- IIb 1 Progression of blastoderm until yolk is covered up by 1/2.
- IIb 2 Progression of blastoderm until yolk is covered up by 3/4.
- II 1 Progression of blastoderm until yolk is covered up by 3/4, blastopore open.
- II 2 Blastopore closed. The head region of the embryo apparent.
- III 1 Tail starts to separate from the yolk. The length of the free tail is smaller or equal than 1/2 the head length.
- III 2 The length of the free tail is greater than 1/2 the head length.
- IV The tail extended 1/4 the length of the yolk sac.
- V The tail extended 1/2 the length of the yolk sac.
- VI The tail length greater than 3/4 of the length of the yolk sac and hatching.

REFERENCES

ALHEIT, J., WAHL, E. and CİHANGİR, B., 1987. Distribution, Abundance, Development rates, Production and Mortality of Sprat Eggs. ICES C.M. 1987 / H:45. 7p.
 CİHANGİR, B., Reproductive biology of European pilchard (*Sardina pilchardus* Walb.) in the Aegean (in prep.).
 MOSER, H.G. and AHLSTROM, E.H., 1985. Staging anchovy eggs. In R. LASKER (editor), An egg production method for estimating spawning biomass of pelagic fish: Application to the Northern anchovy, *Engraulis mordax*. U.S. Dep. Commer., NOAA Tech. Rep. NFMS 36 : 37-41.
 PEREZ, N. and RODRIGUEZ, M., 1988. Histología de los folículos post-ovulatorios de *Sardina pilchardus* (Walb.) de la plataforma Nor-Atlántica de la Península Ibérica. Primeros resultados. Inf. Tec. Inst. Esp. Oceanogr. No 68: 11 p.
 RUSSEL, F. S., 1976. The Eggs and Planktonic Stages of British Marine Fishes. Academic Press, London, 524 p.

