

**Introduction**

The unique adaptability of the green alga *Dunaliella* to grow in a wide range of salt concentrations, has made it a favorite subject for detailed physiological and biochemical investigations, aimed at its utility as a source of some fine chemicals such as  $\beta$ -carotene and glycerol or a source of single cell protein for rearing rotifers and brine shrimp, *Artemia*. It is well known that the concentrations and the ratios of the essential nutritive elements, strongly affect algal growth and photosynthetic rates (FISHER *et al.*, 1981). The main target of this work is to maximize *Dunaliella salina* growth with the most economical culture component supply.

**Material and methods**

The methodology of experimental design, namely, the random balance, the fractional factorial and the steepest ascent designs (SATTERTHWAITE, 1959; COCHRAN and COX, 1957) were used to develop a new synthetic optimized medium for the growth of *Dunaliella salina*. All major cations (Na, K, Mg and Ca) were added as chloride /salts and anions ( $\text{HCO}_3$ ,  $\text{SO}_4$ ,  $\text{NO}_3$  and  $\text{PO}_4$ ) as sodium salts. The trace metals (Zn, Mn, Mo, Co, Cu and Fe) were added in chelated form with EDTA. Vitamin B<sub>12</sub> was added to all media at a level of  $1 \mu\text{g/l}$ . Culture media were inoculated under sterile conditions with actively growing *Dunaliella salina*, adjusting its initial concentration to  $10^4$  cell/l. Experiments were performed in triplicates. Cultures were grown in incubator at light intensity 4 k Lux and temperature of  $25 \pm 1^\circ\text{C}$ . Experiment duration lasted for 9 days. Population density was estimated by cell count on a hemacytometer.

**Results and discussion**

The random balance design was used first to evaluate the main effect of 15 nutritive elements as mentioned previously plus the effect of  $\text{H}_3\text{BO}_3$  at 2 levels of concentrations (+1) and (-1), which were chosen to express the highest and the lowest element concentrations used in the known artificial sea water media (ASP-2, ASP-6, ASP-12, ASM and Muller media). Data gained using this design showed that alga was tolerant to a wide range of macroelement concentration changes. Concerning micro- elements and  $\text{H}_3\text{BO}_3$  the best algal yield was achieved at the levels given in table 2. For optimizing the major cations and anions concentrations in relation to algal yield the  $2^{8-4}$  fractional factorial design was used, where the concentrations of the media on which the alga attained its maximum yield in the previous design was taken as a middle point in defining the (+1) and (-1) levels for this design. Results are given in table 1. After statistical treatment of the data we can conclude that algal yield was only significantly affected by the concentration changes of K, Mg, PO and  $\text{CO}_3$ . For optimizing their concentrations, a set of experiments was done using the steepest ascent method, where the composition of medium No 15 on which the algal yield was maximum (table 1) was taken as original point. The highest mean algal yield ( $18 \times 10^6$  cell/ml) was achieved on growth medium No 8. This is about 9-10 times greater than those recorded in the literature at about the same conditions of cultivation used in our experiments. In conclusion we are recommending a new medium for best *Dunaliella salina* growth as given in table 2.

Table 1.  $2^{8-4}$  fractional factorial design and steepest plan.

Factors	NaCl X <sub>1</sub>	KCl X <sub>2</sub>	MgCl <sub>2</sub> X <sub>3</sub>	CaCl <sub>2</sub> X <sub>4</sub>	Na <sub>2</sub> SO <sub>4</sub> X <sub>5</sub>	NaHCO <sub>3</sub> X <sub>6</sub>	NaNO <sub>3</sub> X <sub>7</sub>	NaH <sub>2</sub> PO <sub>4</sub> X <sub>8</sub>	culture density after 9 days $10^6$ cell/ml
Level mM/l									
-1 level	1000	8	25	8	15	1.5	2	0.05	
+1 level	2000	16	55	18	25	2.5	5	0.10	
0 level	1500	12	40	13	20	2.0	3.5	0.075	
Variation unit ( $\lambda$ )	500	4	15	5	5	0.5	1.5	0.025	
Experiment									
1	-1	-1	-1	-1	-1	-1	-1	-1	4.39
2	+1	-1	-1	-1	+1	+1	+1	-1	3.51
3	-1	+1	-1	-1	+1	+1	-1	+1	6.06
4	+1	+1	-1	-1	-1	-1	+1	+1	4.56
5	-1	-1	+1	-1	+1	-1	+1	+1	6.97
6	+1	-1	+1	-1	-1	+1	-1	+1	4.61
7	-1	+1	+1	-1	-1	+1	+1	-1	4.63
8	+1	+1	+1	-1	+1	-1	-1	-1	3.47
9	-1	-1	-1	+1	-1	+1	+1	+1	0.28
10	+1	-1	-1	+1	+1	-1	-1	+1	5.09
11	-1	+1	-1	+1	+1	-1	-1	-1	4.44
12	+1	+1	-1	+1	-1	-1	+1	+1	4.64
13	-1	-1	+1	+1	+1	+1	-1	-1	4.07
14	+1	-1	+1	+1	-1	-1	-1	-1	4.59
15	-1	+1	+1	-1	-1	-1	+1	+1	7.66
16	+1	+1	+1	+1	+1	+1	+1	+1	5.17
Regr. coef. ( $b_i$ )	-0.23	0.5	0.46	-0.19	0.26	-0.46	-0.41	0.47	
$b_i \lambda$		2.0	6.9			-0.23		0.012	
Initial level	1000	16	55	18	15	1.5	2	0.100	
Level on path.									
1	1000	18	62	18	15	1.25	2	0.112	12.90
2	1000	20	69	18	15	1.00	2	0.124	14.33
3	1000	22	76	18	15	0.75	2	0.136	14.09
4	1000	24	83	18	15	0.50	2	0.148	13.76
5	1000	26	90	18	15	0.25	2	0.160	15.43
6	1000	28	97	18	15	0.00	2	0.172	13.55
7	1000	30	104	18	15	0.00	2	0.184	14.49
8	1000	32	111	18	15	0.00	2	0.169	18.10
9	1000	34	118	18	15	0.00	2	0.208	15.91
10	1000	36	125	18	15	0.00	2	0.220	16.01
11	1000	38	132	18	15	0.00	2	0.232	17.19
12	1000	40	139	18	15	0.00	2	0.244	16.24
13	1000	42	146	18	15	0.00	2	0.256	15.92
14	1000	44	153	18	15	0.00	2	0.268	16.17
15	1000	46	160	18	15	0.00	2	0.280	17.24

**REFERENCES**

COCHRAN W.G. and COX G.M., 1957.- Experimental design. 2nd. edn., Wiley Publication in Applied Statistics, Wiley, New York.  
 FISHER N.S., JONE G.J. and NELSON D.M., 1981.- Effect of copper and zinc on growth, morphology and metabolism of *Asterionella japonica* (Cleve). *J. Exp. Mar. Biol. Ecol.* 51, 37-56.  
 SATTERTHWAITE F.E., 1959.- Random balance experimentation. *Technometrics*, No.2, 111.

Large aggregations of the jellyfish *Rhopilemanomadica* GALIL, 1990 - a new lessepsian migrant into the Mediterranean (GALIL *et al.*, 1990) - have been observed off the Levantine coasts in the past seven years. This jellyfish, which umbrella could reach a size of one meter, inflicts painful stings. The severity of the stings is related to the area affected and depends on individual sensitivity. The symptoms involve a burning sensation in the contact area, swelling and development of fluidfilled blisters that may persist for days and remain as weal marks. Systemic symptoms in the most severe cases may involve high temperature, fatigue and muscular aches.  
 Nets strung along beaches for the protection of bathers were ineffective as tissue fragments, mainly the tentacles of the oral lobes and of the umbrella, that contain nematocysts, passed through and caused a "stinging water" sensation for bathers. Local municipalities reported a decrease in beach attendance during periods of jellyfish swarmings.  
 In 1991 the joint research effort of the Universities of Trieste, Haifa and the National Oceanographic Institute of Israel aimed at the study of the biology, morphology and distribution of *R. nomadica*. This work summarizes the first observations on its nematocysts.  
 The nematocysts were culled from freshly caught jellyfish. Oral tentacles were excised and immersed in distilled water for 24H at  $5^\circ\text{C}$ , then homogenized. The homogenate was centrifuged repeatedly (at 3500 rpm, 15 min.); the supernatant removed, the pellet was re-suspended in distilled water until satisfactory purification. Nematocysts discharge and fixation of SEM samples follow procedures described in AVIAN *et al.* (1991).  
 Our preliminary results attest to the presence of four types of nematocysts in adult specimens of *R. nomadica*.  
 - *Heterotranchous microbasic eurytele*. It has an everted tubule with a well-defined shaft armed with three helicoidal series of spines. Its capsule, 4-6  $\mu\text{m}$  long, is ellipsoidal.  
 - Large *Holotranchous isorhiza* has sub spherical capsules, 8-12  $\mu\text{m}$  long. The everted tubule is armed with three helicoidal series of triangular spines.  
 - *Heterotranchous isorhiza* has capsules ranging from ellipsoidal to truncated cones, 4-5  $\mu\text{m}$  long. The everted tubule is proximally armed with three helicoidal series of flat, lanceolate spines, and distally armed with short, blunt spines.  
 - Small *Holotranchous a-isorhiza*, the smallest of the nematocysts, has an ovoid to sub spherical capsule only 2-3  $\mu\text{m}$  long, with the everted tubule armed with three helicoidal series of little spinules.

Table 2. Mineral composition of medium No. 8.

Conc. ppm/l	NaCl	KCl	MgCl <sub>2</sub>	CaCl <sub>2</sub>	Na <sub>2</sub> SO <sub>4</sub>	NaHCO <sub>3</sub>	NaNO <sub>3</sub>	NaH <sub>2</sub> PO <sub>4</sub>
1000	32	111	18	15	1.5	2	0.10	
Fe	18	15	0.1	0.02				
Zn	1	1						
Mn	1	1						
Mo	1	1						
Co	1	1						
Cu	1	1						
H <sub>3</sub> BO <sub>3</sub>	2	2						
NaH <sub>2</sub> PO <sub>4</sub>	0.150	0.150						
400	0.1	0.02						




Fig. 1. SEM micrographs of the nematocysts of *R. nomadica*. a, discharged *Heterotranchous microbasic eurytele*; b, discharged large *Holotranchous isorhiza*; c, discharged *Heterotranchous isorhiza*; d, discharged small *Holotranchous isorhiza*.

The eurytele type is common both in the tentacles and in the gastric filaments, the large holotranchous isorhiza is more frequent in the scapulate tentacles than in the oral lobes, and the small holotranchous isorhiza is the commonest type, widely distributed in all areas.

It is of interest that a co-generic species, *R. esculenta* Kishinouye, similarly has four nematocysts types in the adult (CHEN & DING, 1981); the classification proposed for the *R. esculenta* nematocysts is otherwise not correspondent with our observations. CHEN & DING's classification is based solely on light microscope observations, and it is possible that their anisorhiza-type nematocysts are in fact the isorhiza-type nematocysts we have identified in *R. nomadica*.

**REFERENCES**

AVIAN M., Del NEGRO P. and ROTTINI SANDRINI L., 1991.- A comparative analysis of nematocysts in *Pelagia noctiluca* and *Rhizostoma pulmo* from the North Adriatic Sea. *Hydrobiologia*, 216/217: 615-621.  
 CHEN J. and DING G., 1981.- On the nematocysts of the various stages of development of *Rhopilema esculenta* Kishinouye (Cnidaria, Scyphozoa). *Acta Zool. Sin.*, 27 (4): 310-317.  
 GALIL B.S., SPANIER E. and FERGUSON W.W., 1990.- The Scyphomedusa of the Mediterranean coast of Israel, including two lessepsian migrants new to the Mediterranean. *Zool. Meded. Leiden*, 64: 95-106.