

The northward flow of Mediterranean water in the Adriatic Sea in the light of oxygen isotope measurements

Barbara STENNI*, Paola NICHETTO** and Antonio LONGINELLI*

*Università di Trieste, Istituto di Mineralogia e Petrografia, TRIESTE (Italia)

**Università di Trieste, Dipartimento di Biologia, TRIESTE (Italia)

Despite the research work carried out by several authors the Adriatic sea is not yet known in detail from the point of view of circulation and distribution of the characterizing variables like salinity, temperature and density of the water. Our knowledge regarding the distribution of the oxygen and hydrogen isotopic composition of Adriatic seawater and the latitudinal and vertical changes of these variables is practically inexistent. To start the collection of reliable data in this field we analyzed up to now a few hundred samples of Adriatic seawater from vertical profiles from southern, central and northern Adriatic (Figure 1). For many of these samples other variables (e.g. salinity, temperature and density) are available and were or will be published by other authors. Our isotopic measurements, though largely incomplete, already yielded data which are of importance for a better knowledge of the circulation in this sea. In fact, sets of bottom water samples between Venice and Trieste showed repeatedly $\delta^{18}\text{O}(\text{H}_2\text{O})$ values ranging from +1,1 to +1,4 per mil (V-SMOW). These are very positive values which drastically differ from those of surface water samples which generally range, in this area, between +0,8 and -0,5 per mil. In this case the depletion in ^{18}O is clearly and obviously due to the inflow of river water whose $\delta^{18}\text{O}$ is considerably negative. The range of values goes from about -12,5 per mil in the case of the Adige river to about -10,0 per mil in the case of the Po river to values of about -8 to -9 per mil for other minor rivers in the area. This means that only when a marked cooling of surface water takes place in this area or strong winds increase the dynamic of water masses, a real homogeneization of the different water layers takes place in northern Adriatic, as shown, e.g., by sets of samples collected in the winter period. When, on the other hand, surface warming and/or the absence of strong wind increase the vertical stability of different water layers, no vertical mixing takes place and even diffusion processes seem to be drastically reduced. Under these conditions, bodies of Mediterranean water from the eastern and central basins -characterized by very positive $\delta^{18}\text{O}$ values- may apparently flow all the way to the northernmost section of the Adriatic sea as a thin bottom layer. Despite a course of hundreds of kilometers this bottom water layer, under favourable conditions, may largely preserve its original isotopic composition through time and space. In fact, eastern and central Mediterranean are the only sources of isotopically positive water which may explain the most positive values obtained. The northward evolution of the $\delta^{18}\text{O}(\text{H}_2\text{O})$ values from the Otranto channel and the dynamic of the Adriatic water masses are discussed in some detail on the basis of the isotopic values obtained and their comparison with other classical variables measured in seawater.

