

**Sedimentological characters of the area North of Eratosthenes seamount  
(Cyprus sector of the Eastern Mediterranean Sea)**

G. CATANI<sup>o</sup>, G. LENARDON<sup>o</sup>, G. TUNIS<sup>o</sup>, A. BONAZZI\* , A. VINCI+ and  
R. BARTOLE#

<sup>o</sup> Institute of Geology, University of TRIESTE (Italy)  
\*Institute of Mineralogy, University of PARMA (Italy)  
+ Institute of Petrography, University of PARMA (Italy)  
# Institute of Geology, University of URBINO (Italy)

Three small basins occur in the sector North of Eratosthenes Seamount and South of Cyprus Island in the Eastern Mediterranean Sea (Fig. 1). The largest one is the Eratosthenes basin which is characterized by compressive and extensional tectonics on its northern and southern flanks respectively (FINETTI and MORELLI, 1973; CATANI *et al.*, 1983; ANASTASAKIS and KELLING, 1991). From a seismo-stratigraphic point of view, high resolution Sparker and 3.5 kHz S.B.P. surveys point out the existence of Messinian evaporitic substratum, dipping northward beneath the south-verging Cyprus Arc thrust system. The overlying Plio-Quaternary sediments transgress on the Messinian substratum with a contact of onlap type. Their attitude is almost everywhere undisturbed, except for the northern part of the basin where they are involved in the compressional tectonics of the most external and youngest part of Cyprus Arc. As for the provenance of the Plio-Quaternary sediments, both core samples and seismic profiles (sparker and S.B.P.) demonstrate that they mainly slid down from the flanks of the Eratosthenes Seamount, by means of prevailing slump and mass-flow mechanisms, these phenomena being clearly recorded by sedimentological data. The remaining terrigenous materials coming from Cyprus are trapped by a morphological high, acting as a barrier; in the western sector of the examined area the sediments sliding down along the slope are trapped in a narrow small and deep basin (near cores 8 and 9 in Fig. 1). Organic fraction emphasizes the existence of two sedimentary domains in this sector. In fact, reworked material, with high interspecific differentiation, characterizes the northern area; conversely, in the South the number of species is low and with very high intraspecific abundance. Grain size analyses also confirm the existence of these two domains; the surficial sediments are fine and homogeneous in the whole area, while the deeper sediments of each core prove that terrigenous input prevails in the cores near Cyprus, while pelagic sedimentation dominates in the southern cores and in the GAN 17 core (Fig. 2). Due to the low contents of smectite, the mineralogical study of clay fraction demonstrates the limited influence of the nilotic sediments in this area, while the high values found for illite and chlorite would mean that the eolian transports played an important role (Fig. 2). Based on the ash layers analysis on ME 4 core, sedimentation rate is very high (24 cm/1000 yrs) South of Cyprus, from 18,000 up to 8,000 years B.P., while it abruptly diminishes to 5 cm/1000 yrs in the last 8,000 years. Such a difference seems related to the sea-level lowstand that occurred during the 20,000-10,000 yrs B.P. interval (STANLEY and MALDONADO, 1979), with a consequent increasing of terrigenous contributes from Cyprus. These data are confirmed by the study of Y-2 and W-3 ash layers recorded in the GAN 17 core. They display a constant low rate of sedimentation (4,1 cm/1000 yrs, from 120,000 to 100,000 yrs B.P.), being far from the mainland and characterized only by pelagic sedimentation. This situation seems well confirmed by heavy mineral composition (epidotes, amphiboles, clinopyroxenes and orthopyroxenes). In fact, it can be observed that a gradual decrease of epidotes (typical mineral of Cyprus suite) occurs from the bottom to the top of the cores, which should be a direct consequence of a similar reduction of the clastic input from Cyprus. Moreover, in the eastern cores an increase of amphiboles and clinopyroxenes and a decrease of epidotes confirm the more important provenance of the sediments from Eratosthenes Seamount. In conclusion, the recent sedimentation study of this sector of the Levantine Sea allowed the identification of three areas with a peculiar prevailing sedimentation mechanism of different material supply : the first, along the southern Cyprus continental slope, is influenced by Cyprus terrigenous input, the second, placed North of Eratosthenes Seamount, is mostly characterized by gravity flows, and lastly, a western area, steadily dominated by pelagic sedimentation. In all the examined areas, nilotic contributes are rather reduced, while the eolian transport influences the northern sector.





Fig. 2



**REFERENCES**

ANASTASAKIS G. and KELLING G., 1991. - Tectonic connection of the Hellenic and Cyprus arcs and related geotectonic elements. *Mar. Geol.*, 97 : 261-277.  
CATANI G., LENARDON G., MARCHETTI A., TUNIS G. and VINCI A., 1983. - Sedimentological and seismic features in the Cyprian sector of the Eastern Mediterranean Sea : preliminary results. *Boll. Oceanol. Teor. Appl.*, 4 : 311-318.  
FINETTI I. and MORELLI C., 1973. - Geophysical exploration of the Mediterranean Sea. *Boll. Geof. Fis. Appl.*, 15 : 263-241.  
STANLEY D.J. and MALDONADO A., 1977. - Nile cone. Late Quaternary stratigraphy and sediment dispersal. *Nature*, 266 : 129-135.