

**Biological responses of mussels transplanted in a heavy metal polluted environment of North Tyrrhenyan Sea**

Francesco REGOLI and Enzo ORLANDO

Dipartimento di Biomedicina Sperimentale, Università di PISA (Italy)

Many studies have indicated that trace metal concentrations in mussels reflect the condition of the environment (review by PHILLIPS, 1980), and this is the reason why they are often chosen as "bioindicators" for monitoring programs.

It has also been outlined the importance of investigations on the biological effects of pollutants which could possibly relate the level of pollution in the environment with the health conditions of the organisms.

However, biological responses to heavy metals (such as accumulation and excretion, cellular and physiological effects...) have been usually studied in laboratory conditions and seldom in field experiments.

In this work, control mussels *Mytilus galloprovincialis*, obtained from a marine farm, were transplanted in plastic cages to a heavy metal polluted site of the North Tyrrhenian Sea (Scarlino), and removed by SCUBA diving at each sampling time.

Analysis of heavy metals revealed a strong accumulation in the digestive gland of transplanted mussels. After 1 week concentrations increased from less than 2 µg/g to 60 µg/g for Pb, from 19 to 100 µg/g for Mn, from 600 to 4000 µg/g for Fe. These high levels of metals in Scarlino are related to the presence of factories which produce sulphuric acid and titanium dioxide from pyrites and titanium minerals.

A "digestive gland index" was calculated as the ratio between digestive gland and shell weight. Variations of this d.g. index reflect changing digestive gland weight, shell being considered a much less variable parameter.

Figure 1 shows "d.g. index" in mussels from the farm and in those transplanted to Scarlino.

During the gametogenesis, the digestive gland is progressively penetrated by gonadal tissues, this determining the increase of digestive gland weight and, consequently, of the "d.g. index" observed in control mussels (Fig. 1A). This is confirmed by the fact that the highest values of the d.g. index were observed just before spawning period, when gonadal development is maximum.

The initial decrease of the "d.g. index" in transplanted mussels (Fig. 1B) indicates a loss of weight by this organ. This fact probably reflects the presence of defence mechanisms, such as a decrease of filtration rate, which reduce the exposure to metals (with obvious consequences on the nutritional state of the organisms), as well as the presence of a general stress induced by the environmental conditions.

The "d.g. index" resulted always lower in transplanted than in control mussel, this indicating the persistence of stress by heavy metals.

Furthermore, low values of the d.g. index in transplanted organisms also indicate a reduced gonadal development as a consequence of a low availability for gametogenesis of energy reserves which are rapidly utilized during stress period (THOMPSON *et al.*, 1974).

Responses to heavy metals were investigated also at the subcellular level where a severe disfunction of the lysosomal compartment was observed (REGOLI, 1992).

Transplanted mussels showed a reduced lysosomal membrane stability, the presence of enlarged secondary lysosomes and an enhanced production of lipofuscin in tertiary lysosomes.

These subcellular alterations as "early warning systems" can reveal a toxicity of pollutants before damages shift to higher levels of biological organization, and they could be used in monitoring programs as early biomarkers of stress.




FIGURE 1

A) Control mussels  
B) Transplanted mussels

**REFERENCES**

- PHILLIPS J. H., 1980. - Quantitative aquatic biological indicators. Applied Science Publishers Ltd., London  
REGOLI F., 1992. - Lysosomal responses as a sensitive stress index in biomonitoring heavy metal pollution. *Mar. Ecol. Prog. Ser.* (n° 2511) in press  
THOMPSON R.J., RATCLIFFE N.A., BAYNE B.L., 1974. - Effects of starvation on structure and function in the digestive gland of the mussel (*Mytilus edulis* L.). *J. Mar. Biol. Ass. U.K.* 54: 699-712