

Environmental conditions which determine chrysophyte *Apedinella* development and blooms

Pia Elena MIHNEA

Romanian Marine Research Institute (Romania)

The modifications of phytoplanktonic community during the last decades (MIHNEA, 1987, MIHNEA *et al.*, 1980) also include the occurrence of some new representatives the number and frequency of which are still increasing. One of them is the chrysophyte *Apedinella* (THRONDSEN, 1971) that produces blooms. Our observations were carried out from 1975 to 1989 in the southern part of the Romanian inshore area along the 13 transects, 5, 10, 20 m isobaths. Samples (1,000 ml) were collected monthly at standard depths (0, 5, 10, 20 m), between February and October. On the whole 2,332 phytoplanktonic samples were analysed. Determinations regarding macronutrients and physical parameters were also done. As macronutrients are not limiting within the studied area, a correlation between this new organism and organic compounds produced by *Skeletonema* was found to help partially the explanation of its bloom (MIHNEA, 1992). Although *Apedinella* was present in Spring-Summer and sometimes in Autumn associations (Table 1), its massive development or bloom could be correlated with the large variation of silicium level (Table 2).

Table 1 : The frequency of *Apedinella* sp. population as its maximum density (no cells/lx10³; underlined values no cells/lx10⁶)

Year/ Month	MARCH	APR.	MAY	JUNE	JULY	AUG.	SEPT.	OCT.
1975	0	<u>3.5</u>	140.0	33.0	0	0	0	0
1976	0.2	<u>14.5</u>	5.2	8.4	0.2	0	0	10.6
1977	0	1.4	0	0	0	2.1	1.5	0
1978	0	<u>3.4</u>	0	1.7	11.8	36.4	0	0
1979	108.0	<u>132.0</u>	1.5	31.5	0	0	0	3.0
1980	62.4	8.1	0	1.3	0	1.0	0	1.0
1981	0	0	15.8	0	0	0	0	0
1983	0	<u>1.0</u>	<u>21.5</u>	23.0	41.6	915.0	0	7.6
1984	0	<u>23.4</u>	<u>1.0</u>	237.0	4.0	14.0	0	0
1985	0	34.0	2.0	12.0	10.0	15.0	0	0
1986	0	6.0	<u>13.3</u>	38.0	6.0	12.0	6.0	0
1987	0	16.8	<u>151.0</u>	15.7	0	12.6	0	0
1989	49.0	0	0	2.1	14.9	12.8	0	17.0

Table 2 : Silicium level during the whole year within the studied area (CUINGIOGLU, unpubl.)

Year	Total No of samples	Class size/No of samples(N)/ range of Si-SiO ₃ variation	
		> 1000 µg/l	< 1000 µg/l
1983	349	41	308
		1006-2000	0-960
1984	356	37	319
		1017-3740	0-904
1985	335	15	320
		1011-4960	53-845
1986	260	25	235
		1005-2685	78-945
1987	231	69	162
		1005-6125	105-990

April and May are the months when this chrysophyte blooms and correspond with the most reduced level of Si-SiO₃. If we only consider the 1986 results, when *Apedinella* cell no/l reached respectively 21.5 and 13.3 x 10⁶, silicium concentrations varied on one hand from 0 to 400 µg/l in 35.54% and from 400 to 800 µg/l in 53.85% of the samples (N=104), and on the other hand from 0 to 400 µg/l in 54.0% and from 400 to 800 µg/l in 36.78% of the samples (N=87).

The silicium depletion is due to *Skeletonema* and some other co-associate diatoms which previously developed for a long while; this event plays an important role in shifting the species composition from diatoms to algal flagellates supposed to have limited or to have not Si requests. The sequence is very short but *Apedinella* "takes advantage" and begins its intense divisions. Some other factors could be involved, e.g. the total radiant energy. An increase in total radiant energy during the last decade of April and the first decade of May (Table 3) also could be connected to the promoting of an accelerated division rate of *Apedinella* if it is combined with a 6-10°C range of sea temperature (MIHNEA, 1992). The superimposing of : *Skeletonema* bloom, decrease of Si, 6-10°C range of sea temperature and a rapid increase of light (more than 1000 cal cm⁻²) should be considered the "conditioning" of sea water for the described chrysophyte development.

During 1984, 1985 and 1987 the total radiant energy had increased in first decade of May, temperature was 10-16°C and the silicium level higher than values determined in 1983 and 1986's Spring; in such a "scenario" the bloom of chrysophyte *Apedinella* didn't take place. We cannot omit the concurrence between species as well the physical forces that could be other conditions of mass development. A tight concurrence between *Apedinella* and dinophyte *Heterocapsa triquetra* was observed, too; they both compete for some "excreted substances", or maybe try to impose their own association to medium diatoms which cannot develop in.

If *Apedinella* cannot develop strongly enough in April, windy Spring weather could stop it by dispersion of its patches.

In conclusion, in eutrophicated marine ecosystems, individual species are controlled by a large set of environmental factors. If anyone changes significantly, it upsets the existing balance and induces a new ecological equilibrium. The bloom is caused by the coupling of the physiological capabilities with the environmental conditions.

ACKNOWLEDGEMENTS

I am deeply indebted to my colleague E. CUINGIOGLU for providing useful unpublished data on silicium.

REFERENCES

- MIHNEA P.E., 1987.- The eutrophication process in the inshore Romanian Black Sea area. *Rev. Roum. Biol. - Biol. Anim.* 32, (2) : 149-155.
 MIHNEA P.E., 1992.- Blooms of *Chrysophyta* in the Black Sea. *Proceed. of Fifth Intern. Conf. on Toxic marine phytoplankton*, Newport, 1991 (in press).
 MIHNEA P.E., CUINGIOGLU E. et BILAL I., 1980.- Modifications des communautés phytoplanktoniques littorales sous l'influence du phénomène de pollution. *Ves Journées Etud. Pollution*, Cagliari, CIESM : 869-876.
 THRONDSEN J., 1971.- *Apedinella* gen. nov. and the fine structure of *A. spinifera* (Thronsd.) comb. nov. *Norw. J. Bot.*, 18, (1) : 48-64.

Rapp. Comm. int. Mer Médit., 33, (1992).