

Dubravka REGNER

Institute of Oceanography and Fisheries, SPLIT (Croatia)

The aim of this paper is to determine the consequences of progressive eutrophication effects caused by human activity on the composition and density of copepods.

Zooplankton material for this study was collected from bottom to surface at five stations in the eastern Adriatic with HENSEN plankton net (73/100, silk N° 3). One of them was in the coastal waters in front of Dubrovnik (Gruz) harbour, two in the Vela Luka Bay (channel region of the middle Adriatic) and the last two in the open middle Adriatic (Fig. 1).




Fig.1 The study area

Fifteen years of investigations (1977-1991) on copepods in front of Dubrovnik harbour have shown that about 30 species are present in summer (REGNER, 1986, 1987, 1989, 1991). Although some predominantly pelagic copepods such as : *Neocalanus gracilis*, *Calocalanus plumulosus*, *Clausocalanus parapergens*, *Pleuromamma gracilis*, *Heterorhabdus papilliger*, *Corycaeus flaccus* and some others appear from time to time, neritic species usually occur with higher densities. *Acartia clausi* is always dominant in summer, with an extraordinary high percentage of up to 80 %. In 1991, the percentage of *Acartia clausi* was 93% of all other copepods. In addition recent results (July 1991) indicate that the density of copepods show an increasing trend of about 1300 copepods/m³. The increasing trend of the percentage of *Acartia clausi* and the increasing copepod density we can connect with the progressive eutrophication of eastern Adriatic coastal waters and the increasing phytoplankton density in fifteen years period.

In Vela Luka Bay, about 35 species of copepods were found in 1990. The maximum number of species was found at the entrance to the bay, exposed to the strongest impact of open sea waters. Nevertheless, the species composition is diverse in the inner part of the bay, with some pelagic species in addition to those of neritic origin. So, the presence of the copepods: *Lucicutia flavicornis*, *Mecynocera clausi*, *Euchaeta hebes*, *Heterorhabdus papilliger* and some others, predominantly pelagic species, indicate a mixing of coastal and pelagic influences throughout the study area. Dominant copepods were: *Acartia clausi* constituting as much as 93% in July, *Centropages kroyeri* (up to 50% in autumn), *Temora stylifera*, *Labidocera wollastoni* etc. Apart from the high percentage of *Acartia clausi* in summer, the eutrophication effects of human activities can be seen in the inner part due to an unexpected high biomass for such a shallow station. This phenomenon was observed especially in summer.

At stations close to the Jabuka Pit and Palagruza (open waters of the middle Adriatic), about 50 species of copepods were found. Although this number is smaller than expected, we can explain it by seasonal samplings of zooplankton in 1986 and 1987. *Ctenocalanus vanus*, the most wide-spread Adriatic species was dominant of the present copepods with about 11%. Furthermore *Acartia clausi* was present by 6%, *Clausocalanus jobei* by 4.6%, *Centropages typicus* by 4.4% etc. From time to time, with just a few specimens : *Clausocalanus mastigophorus*, *Calocalanus plumulosus*, *Clausocalanus parapergens*, *Haloptilus longicornis*, *Lucicutia flavicornis*, *Macrosetella gracilis*, *Scolecithricella dentata*, *Corycaeus flaccus* and some other predominantly pelagic species appeared, as a consequence of dynamics of water masses and mixing of coastal and open-sea waters at the studied stations. The number of copepods per m³ is low-about 112/m³ which is usual for open waters of the middle Adriatic.

Conclusions

From the results mentioned above we conclude:

1. The recent results in front of Dubrovnik (coastal sea) have shown progressive changes under the influence of eutrophication, such as the increasing trend of the percentages of *Acartia clausi* and the increasing copepod density.
2. In the Vela Luka Bay (channel region) the eutrophication effects of human activities can be seen in the upper part of the bay through unexpectedly high biomass and high percentage of *Acartia clausi*, too.
3. No changes have been recorded yet from the open middle Adriatic.

REFERENCES

- REGNER D., 1986.- Eutrophication effect on the copepod community of the eastern Adriatic coast. *Rapp. Comm. int. Mer. Médit.*, 30, 2, p. 202.
- REGNER D., 1987.- The impact of pollution on the copepod community from the eastern Adriatic coast. *Chemosphere*, 16, (2/3): 369-379.
- REGNER D., 1989.- Copepod communities in the middle Adriatic as affected by different levels of pollution. *Toxicological and Environmental Chemistry*, 20/21: 217-225.
- REGNER D., 1991.- The progressive changes of the copepod community from the eastern Adriatic coast caused by eutrophication. *Toxicological and Environmental Chemistry*, 31/32 : 433-439.