

*Istituto per la Pesca Marittima, ANCONA (Italia)

**Istituto delle Grandi Masse, VENEZIA (Italia) (permanent affiliation: Institute of Oceanography and Fisheries, SPLIT, Croatia)

***Osservatorio Geofisico Sperimentale, TRIESTE (Italia)

****Hydrographic Institute, SPLIT (Croatia)

The Adriatic Sea basin-wide circulation and its temporal variability is reviewed based on the results of analysis of hydrographic data collected during POEM cruises. Major well known characteristics of the circulation are revealed from the present data set which covers the period from October 1985 to April 1987. A prominent signal associated with the seasonal variability, is identified the outflowing water along the Italian coastline. Differences between autumn and spring in the vein of cold and fresh water along the Italian shelf break, manifest mainly in the temperature field. In the surface layer during the stratified season in the Southern Adriatic the fresh water spreads over the entire surface. On the other hand, during the spring i.e. the period when the sea is vertically homogeneous, the fresh water remains confined in the surface alongshore boundary layer over the entire length of the Italian shoreline. Layers below the seasonal thermocline at the eastern portion of the sea display very weak seasonal signal. A strong signal associated with the inter-annual variations have also been documented from the analysed data set. It mainly appears in the salinity field. It is shown that in the spring 1986 the salinity averaged over the entire water column north of the Palagruza Sill is by 0.3 PSU smaller than in the spring 1987. The same but less prominent difference is noted in the southern Adriatic. These differences are related to the variations in climatic conditions over the area, the river run-off and the Mediterranean water inflow.

Gordana BEG and Miroslav GACIC

Institute of Oceanography and Fisheries, SPLIT (Croatia)

Kastela Bay is a small semienclosed basin with total area of 61 km² and average depth of 23 m. The bay communicates with the adjacent sea through a relatively wide (1.8 km) and deep mouth (mean depth about 40 m). The principal freshwater source is Jadro River with relatively small discharge (yearly mean about 10 m³ s⁻¹).

Seven current measurement experiments of approximately two month duration were made in the Kastela Bay from 1980 to 1990. Current measurement stations were placed in the center and in the mouth of the bay where water exchange with the adjacent sea takes place. In each experiment measurements were made simultaneously at more than one station and several depths. This made possible the determination of vertical and horizontal distribution of current field. Measurements in various seasons allows us to detect intra-seasonal differences in the current field.

Current meter data were analyzed together with appropriate hourly wind and atmospheric pressure data recorded at the meteorological station in the vicinity in order to study the effects of the atmospheric synoptical disturbances on the current field and water exchange with the adjacent sea. Currents and wind stress vectors were decomposed into the east-west and north-south components. All these time series were then low-pass filtered with 24m214 filter (THOMPSON, 1983) to reject tidal and other high frequency motions. Besides of the detailed analyses of low-pass filtered time series, correlation coefficients between north and east components of wind stress and currents were computed for periods with wind speed greater than 5 m/s in order to study the momentum balance (WINANT, 1980). Basic statistical characteristics (mean, standard deviations) were also computed. Correlation coefficients between current components and atmospheric pressure were computed for entire time series. Using linear interpolation for the current profiles, water transports and renewal times were obtained for stations in the mouth of the bay. Schemas of water transports were given for strong wind events.

The most obvious characteristic of all time series is strong variability on a weekly time scale.

Current meter data in the mouth of the bay show well defined two-layer structure for the periods with strong wind events, with surface current in phase with wind, while in deeper layers the phase-lag between wind and currents is 180 (GACIC, 1982; GACIC *et al.*, 1987). Correlation coefficients between currents and wind stress components, as well as average vectors give frictional momentum balance. Great negative correlation coefficients between the current components show polarization of the currents in the mouth in the east-west direction during the strong wind events. Correlation coefficients between east component of currents in the surface layer and atmospheric pressure show incoming flow during the periods with low pressure and outgoing flow during the periods with high pressure. The motions in the lower layers are in the opposite directions. Low pressure is often connected with scirocco (SE) wind, that also drives incoming flow in the surface layer and high pressure is connected with bora (NE) wind that drives outflow in the surface layer (ZORE-ARMANDA, 1980). When the scirocco (bora) wind stops the pressure starts to raise (fall) and opposite circulation in the mouth is established. Deviations from these typical synoptical situations were also observed.

Magnitudes of currents and correlation coefficients between currents and wind stress and currents and atmospheric pressure were smaller for the stations in the center of the basin. The correlation coefficients between wind stress and current components, as well as average current vectors reveal Ekman momentum balance with Ekman transport in the surface layer and compensatory transport in deeper layers.

Calculations of the water transports and renewal times show that the wind is effective mechanism for water exchange in the bay. Water renewal times are of order of two days during the strong wind events.

REFERENCES

- GACIC M., 1982.- Notes of characteristic of the response of near-shore current field to the onshore wind. *Notes*, No. 47, Institute of Oceanography and Fisheries, Split, 1-6.
GACIC M., DADIC V., KRSTULOVIC N., MARASOVIC I., MOROVIC M., PUCHER-PETROVIC T. & SVILICIC N., 1987.- Near-shore transport processes induced by the wind. *Estuarine, Coastal and Shelf Sci.*, Vol. 24., 35-46.
THOMPSON, R., 1983.- Low-pass filters to suppress inertial and tidal frequencies. *J. of Phys Oceanogr.*, Vol.13., 1077-1083.
WINANT C.D., 1980.- Coastal circulation and wind-induced currents. *Ann. Rev. Fluid Mech.*, Vol. 12., 271-301.
ZORE-ARMANDA M., 1980.- Some dynamical and hydrographic properties of the Kastela Bay. *Acta Adriat.*, Vol. 21.(2), 55-74.