

INPUTS OF PHOSPHORUS AND NITROGEN INTO THE MEDITERRANEAN SEA BY THE RHONE RIVER. VARIABILITY DURING THE LAST 20 YEARS

B. COSTE and P. BÉTHOUX

Centre d'Océanologie de Marseille, Fac. des Sciences de Luminy, 13288 Marseille, France

Continental inputs of phosphorus and nitrogen into the Mediterranean Sea are connected to several features :

- 50% of the primary productivity in the dilution area (Gulf of Lions) are supported by phosphate and nitrate inputs by the Rhone river (COSTE, 1974).

- the continental inputs explain, for a part, the concentrations of phosphate and nitrate in the DEEP WATER OF THE MEDITERRANEAN SEA (MC GILL, 1968; BÉTHOUX AND COPIN-MONTÉGUT, 1986; COSTE *et al.*, 1988). Indeed, input (Atlantic water and river water) and output (Mediterranean Sea water) on the straits of Gibraltar are equilibrated. Modifications in the amounts of phosphorus and nitrogen by river discharges could contribute to modifications in the deep water concentrations. In this paper, we want to present Rhone river data obtained since 1968 and, more particularly, some recent data (1989 and 1990) to discuss their temporal variability.

Water input. The Rhône river is the main contributor to river water input into the western mediterranean basin. Its mean flow is 1700 m³.s⁻¹; Ebro river is 200 m³.s⁻¹, Arno river 103 m³.s⁻¹, Tiber river 234 m³.s⁻¹ (MARTIN and SALIOT, 1992). The Rhône river flow is very irregular: it varies from 500 to 13000 m³.s⁻¹ and it presents a seasonal variability (maxima in autumn and winter and minima in summer) as well as an interannual variability (Table 1).


Table 1. main data concerning the Rhône river flow and nutrient concentrations during the 4 years studied.

Year	1968	1984	1989	1990
Mean annual flow (m3s-1)	1 829	1 678	1 061	1 301
Nutrient inputs (tonnes.y-1)				
N	54 952	53 458	48 591	56 854
P	4 064	6 454	4 712	5 077
Mean annual concentrations (ml-1)				
N	68	72	104	98
P	2,2	3,8	4,4	3,9

Nutrient concentrations. Along the year, the nutrient concentrations vary within a ratio factor lower than 1 to 4. Nitrate are between 50 and 160 µM and phosphate between 2 to 9. These results are in good agreement with those reported by MEYBECK (1982). In 1984, we measured different forms of nitrogen compounds (fig. 1). Nitrite concentrations vary all along the year from 1 to 5 µM, ammonium plus dissolved organic nitrogen (DON) from 20 to 60 µM and nitrate from 50 to 160 µM. Some separated measurements for ammonium and DON show that DON is about 50 to 70% of the total. Then nitrate is the main source of nitrogen input but ammonium and DON have to be taken into account in estimating a mediterranean nitrogen budget.

Seasonal variability in the nutrient concentrations. Maxima concentrations in winter (November to March) and minima in summer (June to September) appear to be a general rule for nutrients (COSTE, 1974). For instance, nitrate concentrations show such a scheme in 1984 (fig. 1). Some exceptions, due to perturbations in the flow of the Rhone river, can be encountered. Such a seasonal cycle does not appear for the other forms of nitrogen. Nitrite does not show a significant seasonality. Ammonium+DON present only a slight seasonal signal with weak maxima in summer and autumn.

Figure 1: Variations in the concentrations of the different forms of nitrogen compounds along 1984.



Seasonal variability in the nutrient inputs. We can compute the nutrient inputs (tonnes.year⁻¹) from the values of the flow and those of the nutrient concentrations. The obtained results are characterized by a high seasonal signal because the higher concentrations values are concomitant to those of the flow. Thus the nutrient inputs are maxima at the beginning of the year and minima in summer with a ratio of 1 to 10.

Interannual variability. Table 1 presents the main features of flow and nutrient input for 4 years. It shows that: 1/ annual nitrate input is rather constant since 1968. Deviations from the mean value are less than 5%. Phosphate annual input is more variable and the recent data are 20% higher than the 1968 data. 2/ the mean concentrations of these nutrients are 50 to 70% higher in 1989 and 1990 than in 1968.

These results lead us to conclude that :

- a high seasonal signal characterizes the nutrient inputs all along the last 20 years. It is due to simultaneity of higher concentrations and higher flow values.

- the annual nitrate input has not varied significantly since 1968. Such a result has to be compared to variability of the mean annual values of the flow. The more recent data of the flow are 30% lower than the earlier. It seems that the low variability of the nitrate input could be explained by the flow variability.

- the variations of phosphate input could be explained by the flow variability but also by an increase in anthropogenic activity.

REFERENCES

- BÉTHOUX J.P. & COPIN-MONTÉGUT G., 1986. Biological fixation of atmospheric nitrogen in the Mediterranean Sea. *Limnol. Oceanogr.*, 31: 1353-1358.
 COSTE B., 1974. Rôle des apports nutritifs minéraux rhodaniens sur la production organique des eaux du Golfe du Lion. *Téthys*, 6: 727-740.
 COSTE B., LE CORRE P. & MINAS H.J., 1988. Re-evaluation of the nutrient exchanges in the Strait of Gibraltar. *Deep-Sea Res.*, 35: 767-775.
 MARTIN J.M. & SALIOT A., 1992. Bilan des apports fluviaux et atmosphériques d'éléments et composés chimiques en Méditerranée occidentale, pp. 61-66, in : 3èmes Rencontres de l'Agence Régionale pour l'Environnement Provence-Alpes-Côte d'Azur, 24-27 Septembre 1991.
 MCGILL D.A., 1969. A budget for dissolved nutrient salts in the Mediterranean Sea. *Cah. océanogr.*, 21: 543-554.
 MEYBECK M., 1982. Carbon, nitrogen, and phosphorus transport by world rivers. *Amer. J. Sci.*, 282: 401-450.

GEOCHEMICAL, SEDIMENTOLOGICAL AND MINERALOGICAL DATA SEDIMENTS OF THE NORTHERN AND CENTRAL ADRIATIC : A MULTIVARIATE STATISTICAL ANALYSIS

A. DE LAZZARI, M. TOSCANO, G. RAMPAZZO AND C. TURETTA

Department of Environmental Sciences, Geo-Mineralogical Section, University of Venice, Calle Larga S.Marta 2137, 30123 Venice, Italy

Several studies of the Adriatic Sea shelf have already been carried out on recent and ancient sediments. The present interest, regarding the study of sediments, is connected with the cognitive research and with the environment protection, which needs, in the complexity of the situation, a natural solution, efficacious and rational. The work, shortly described in this note, has been conducted on recent superficial sediments (thickness from 0-5 cm) of the Adriatic Sea. Sediments samples, for a total of 33, have been drawn in the Summer 1990, along 7 transversal transects in 33 stations in international waters of the northern Adriatic Sea, between the Lagoon of Venice and the junction Gargano-Tremiti-Lagosta.

The following investigations were conducted on the collected materials:

- geochemical : determination of atomic absorption spectrometry of the major (Si, Al, Fe, Mg, Ca, Na e K), minor (Ti, P e Mn) and trace (Zn, Pb, Cd, Cu, Cr, Ni, Co e V) elements;

- mineralogical : definition of the essential qualitative mineralogical composition (light and heavy minerals) by x-ray diffraction analysis, and the study by microscope of some representative samples. The quantity of carbonates was determined by gas-volumetric method.

- sedimentological : determination of the granulometric distribution in the three fractions (sand, silt, clay), and S.S.A. (specific surface area).

A comparative examination of the results obtained by the various investigations has followed, even through a descriptive and multivariate statistical elaboration, which permitted to obtain informations and important results and has pointed out, among the variables, many correlations often difficult to be recognized, operating separately. Experimental data obtained from the various investigations, together with the values of some organic pollutants (PCB, PAH, DDT) associated to particulate input of the several rivers (Po and Adige in particular) in the studied area, have been elaborated by factor analysis (Q-Mode) and analysis of clusters.

The statistical elaboration has allowed to

- identify and define the relationship among determined variables,

- verify the type of connection among particle-size analysis, trace metals and pollutants of organic origin,

- evidence "new connection" among variables (Cd connected to the organic substance; PAH, PCB, DDT, connected to the organic substance and to K),

- synthetize and compare the results obtained with the distribution drawn applying the most classic method of classification. The application of the multivariate statistical analysis, has been preceded by a preliminary analysis of data and of their distribution kind through particular graphic presentation and/or semi-graphic (box-and-whisker plot; stem and leaf display) and the use of non-parametric and robust tests.

The present work must be seen as the starting point of a wider and detailed work of the analysed area.