

ANIMAL SPECIES THREATENED IN THE MEDITERRANEAN ISLANDS : A PROJECT TO PRESERVE VULNERABLE HABITATS

Bruno MASSA

Istituto di Entomologia Agraria, Viale delle Scienze 13, 90128 Palermo, Italy

Mediterranean islands in the last years are facing a serious threat for coastland conservation: the widespread of tourism industry. Following the World Travel and Tourism Council (1992), tourism today has become the most important civil industry in the world : in 1993 it was expected to generate about 3.5 trillions of dollars of world output, that is 6% of the world gross national product, and to double by the year 2005. Tourism is a bigger industry than the auto, steel, electronics or agricultural industries; it grew by more than 57% in the past decade and is expected to grow by 50% before 2000. Some mediterranean islands (e.g. Crete, Rhodes, Cyprus, etc.) in the spring-summer period (May-September) hold almost 10 millions of international travellers, with understandable consequences for the coastline and beach conservation; true, the new economy has given impetus to the agriculture and has generally changed the life style of many people who live in the coast villages, but at the same time has worsened the status of animal and vegetal coenoses through the excessive removal of natural biotopes along coastlines, the building of tourist harbours, the progressive settling of breakwaters, the beach replenishment and the pollution of coasts, which interfere with coastal currents, natural sand sedimentation, as well as terrestrial zoocenoses and phytocenoses.

It is necessary to plan a general project to identify the most important naturalistic areas along island coasts and to prepare an agreement (at least shared by the EC countries) in order to control and improve this threat, without loss for the economy. Mediterranean islands are characterized by harsh landscapes painted with strong colours, superb mountain scenery, woodland, fascinating coastlines and beaches, as well as places of great historical and archeological interest; they are very much suitable to different kinds of nature tourism. Ecotourism is defined as responsible travel to natural areas that conserves the environment and sustains the well-being of local people (BLANGY and EPLER WOOD, 1993). Nature tourism in 1989 generated approximately 7% of all international travel expenditures (CEBALLOS-LASCURAIN, 1993); potential benefits of ecotourism are generation of funds for protected areas, creation of jobs for people who live near protected areas, and promotion of environmental education and conservation awareness (BOO, 1993).

An exemplary case of ecotourism in Mediterranean islands is that reported by PITTET (1994) for Formentera (Balears Is.); tourists conscious of the requirements of the environment and who adapt their behaviour to them contribute to the welfare of local people.

We need to inventory all the preserved coast areas of Mediterranean islands, to analyse speedily their biological-conservationist value using as ecological indicators well-known taxonomic groups (BRICHETTI and MASSA, this volume), and to propose to local governments alternative projects, which consent at the same time ecotouristic exploitation, economic growth and conservation of natural resources. Because much of the threatened biodiversity can be found in comparatively small areas, protection of these areas would ensure the survival of a high number of species, both animals and plants. Analysis of breeding bird distribution, of sites of concentrating migrators and of wintering habitats have revealed that they are good indicators of these key areas.

I conclude by proposing to the scientific community to contribute to this international inventory, and I recall that natural resources are a very important and unique tourist attraction; to do nothing in order to improve their degradation or to contribute to their destruction is a cultural crime like destroying a sculpture made by Michelangelo, as NORTON (1986) wrote: "the damage done when a species becomes extinct is analogous to the damage done when a great work of art is destroyed".

REFERENCES

- BLANGY S. and EPLER WOOD M., 1993. Developing and implementing ecotourism guidelines for wildlands and neighboring communities. Pp. 32-54 in LINDBERG K. and HAWKINS D.E. (eds.), Ecotourism, a guide for planners and managers. Ecotourism Soc., Vermont, USA.
BOO E., 1993. Ecotourism planning for protected areas. Pp. 15-31 in LINDBERG K. and HAWKINS D.E. (eds.), Ecotourism, a guide for planners and managers. Ecotourism Soc., Vermont, USA.
CEBALLOS-LASCURAIN H., 1993. Ecotourism as a worldwide phenomenon. Pp. 12-14 in LINDBERG K. and HAWKINS D.E. (eds.), Ecotourism, a guide for planners and managers. Ecotourism Soc., Vermont, USA.
NORTON B.G., 1986. The preservation of species. The value of biological diversity. Princeton Univ. Press.
PITTET E., 1994. Ecotourism in Formentera, Archipelago of the Balears. *Insula*, 3(1): 48-50.
World Travel and Tourism Council, 1992. The WTTC Report : travel and tourism in the world economy. Bruxelles

THREATENED HABITATS AS A CRITERION FOR SELECTING COASTAL PROTECTED AREAS IN THE MALTESE ISLANDS

Patrick J. SCHEMBRI

Department of Biology, University of Malta, Msida, Malta

Recognising that the sea is one of the Maltese Islands' main resources, recent environmental protection legislation pays particular attention to the coastal zone and shallow seas off the islands. Under this legislation, the terrestrial areas of three islands, and a number of coastal sites, have been declared nature reserves. However, at present there are no marine protected areas within Maltese territory. Marine activities are restricted in certain sea areas round the Maltese Islands, but this is for reasons of security. One factor hindering the establishment of marine protected areas is lack of knowledge as to which ecosystems are in need of protection. In order to address this deficiency, a survey was carried out to identify coastal and shallow water habitats which are threatened. The results of the survey are summarised below. For each habitat type recognised as in need of protection, status, exploitation and threats are reported upon in that order.

Mediolittoral bioconstructions

Status: Platforms formed by the alga *Neogoniolithon notarisii* and the vermetid *Dendropoma petraeum* are common on all gently sloping rocky shores. *Lithophyllum* cushions are only known from a few shores (for example, Xlendi, Gozo and Ghar Lapsi, Malta). These rocky shore bioconstructions are considered vulnerable on a Mediterranean scale (UNEP/IUCN/GIS POSIDONIE, 1990).

Exploitation: No direct exploitation.

Threats: The same as all local rocky shores, mainly from development.

Sea-grass (*Cymodocea nodosa* and *Posidonia oceanica*) meadows

Status: Although common and widespread round the Maltese Islands, in some areas, especially in enclosed or semi-enclosed bays receiving a variety of effluents and subject to certain activities, these meadows have regressed and eroded away, leaving in their place much impoverished thanatocenoses.

Exploitation: Not exploited directly, except for certain types of fishing, but their high productivity makes them one of the most important local sublittoral community types, as in the rest of the Mediterranean (ROS *et al.*, 1985).

Threats: The main threats are: dredging, which causes both mechanical damage and also increases water turbidity and the rate of sedimentation; terrestrial run-off in enclosed bays, particularly that containing sediment and agricultural run-off; cooling water from the local power stations; the hypersaline discharge from reverse osmosis plants; nutrient-rich effluent from sewage; waste from fish-farms; bottom trawling; the use of heavy anchors, which physically damage the meadows; and the illegal use of explosives for fishing. Coastal developments have changed the current and sedimentary regimes in some areas (for example, Marsaxlokk Bay, Malta) and have caused regression of sea-grass meadows.

Posidonia "barrier reefs"

Status: Only very few such formations are known locally. The best documented are those in Mellieha Bay (BORG and SCHEMBRI, 1993) and in Salina Bay, both in Malta.

Exploitation: Not exploited directly, however, these "reefs" are very important in protecting the shore from wave action by absorbing the energy of waves.

Threats: The same as sea-grass meadows in general, but particularly susceptible to mechanical damage, as for example, from boat anchors and moorings.

Halophila stipulacea meadows

Status: Meadows of this Lessepsian immigrant are only known from two Maltese localities, the inner reaches of Marsaxlokk Bay, Malta (LANFRANCO, 1970) and Mgar Harbour in Gozo. The populations at Marsaxlokk have regressed in recent years.

Exploitation: Not exploited.

Threats: Dredging works in connection with the new power station at Delimara, together with pollution resulting from the fishing harbour at Marsaxlokk, have caused a severe decline in the *Halophila* meadows growing in Marsaxlokk Bay.

Deep-water *Cystoseira* communities

Status: Deep-water *Cystoseira* communities based on *C. spinosa*, *C. dubia* and *C. zoeteriodes*, are rare and poorly known in the Maltese Islands, however, it is suspected that they may be threatened, as in other parts of the Mediterranean (UNEP/IUCN/GIS POSIDONIE, 1990).

Exploitation: Not exploited directly, except for some types of fishing.

Threats: Most species of *Cystoseira* are sensitive to pollution, particularly to high phosphate levels and upper infralittoral communities are disappearing from some areas receiving organic pollution; deep-water communities may be similarly affected. Other threats include changes in sedimentary and current regimes due to coastal developments, dumping, and fishing with explosives.

Cladocora cespitosa banks

Status: In Malta this coral forms banks some 20 cm across. These were previously common but are now rare. Large and well-developed banks are particularly rare.

Exploitation: Collected for their curiosity value and for use as decorations in aquaria.

Threats: Overcollecting and mechanical damage from the use of heavy anchors and fishing gear; also, illegal fishing with explosives.

Maërl communities

Status: Apparently rare in the Maltese Islands, although this could be because these communities occur mainly in deep water. It is suspected that those occurring close to the transition zone between the lower infralittoral and the upper circalittoral may be threatened.

Exploitation: Not exploited directly, except for some types of fishing.

Threats: The main threat seems to be from bottom trawling although changes in the sedimentary regime due to coastal development may pose an additional threat in some areas.

Corallegene communities

Status: Occur in deep water (circalittoral) and poorly known locally. It is suspected that those occurring close to the transition zone between the lower infralittoral and the upper circalittoral may be threatened.

Exploitation and threats: The same as for maërl communities.

Caves

Status: Common in Maltese waters and different types exist, ranging from those close to the surface and open to the air, to deep grottoes and tunnels.

Exploitation: Not exploited directly, except for "sight-seeing" by tourist divers.

Threats: The main threat is from divers who enter the caves. These cause both mechanical damage to erect sessile forms, and death of the biota on the ceiling due to air bubbles from diving cylinders becoming trapped there.

REFERENCES

- BORG, J.A. and SCHEMBRI, P.J., 1993. Report on a survey of the marine benthic communities of Mellieha Bay (northern Malta). Malta. Malta Coastal Environment Research Project (CERP). 54 pp.
LANFRANCO, E., 1970. Occurrence of *Halophila stipulacea* (Forsk.) Ascherson (Family: Hydrocharitaceae) in Maltese waters. *The Maltese Naturalist*, Valletta 1(1): 16-17.
UNEP/IUCN/GIS POSIDONIE 1990. Livre rouge "Gérard Vuignier" des végétaux, peuplements et paysages marins menacés de Méditerranée. *MAP Technical Reports Series* 43 : 1-250.
ROS, J.D.; ROMERO, J.; BALLESTEROS, E. and GILI, J.M., 1985. Diving in blue water: the benthos. In : MARGALEF, R. [ed.] Key environments - the western Mediterranean, pp.233-295. Oxford: Pergamon.