

## MARINE ENVIRONMENTAL DATA BASE OF THE ADRIATIC SEA (MAIN RESULTS FROM TIME SERIES ANALYSIS)

Vlado DADIC and Branka GRBEC  
Institute of Oceanography and Fisheries, Split, Croatia

A hydrographic material collected from the Adriatic (ZORE-ARMANDA *et al.*, 1991) by different institutes between 1911 and 1993 is presented. It is stored in MEDAS (Marine Environmental Database of Adriatic Sea) in Institute of Oceanography and Fisheries. Database has been designed as relational using ORACLE RDBMS, and consists of data of climatology, physical oceanography, chemistry, biology, fisheries and pollutants. Data related to air temperature, evaporation, sea temperature and salinity were statistically processed and analysed. Constant fluctuation trends (increase and decrease) for most of the parameters indicate human impact on the sea water properties both in the coastal area and in the open sea. Output data are presented in GIS (Geographical information System) form.

The description of hydrographic properties is primarily based on the data from permanent station at the Split-Gargano transect located along the Palagruza pit and in this region the impacts of both the northern and southern Adriatic waters are felt. Data of air temperature, pressure, and precipitation from some permanent meteorological station (station Trieste, Split and Dubrovnik) were used to interpret the long-term fluctuations of sea temperature, salinity and evaporations both in the coastal area and in the open sea. Main trend analyses are presented in Fig. 1. Long-term variations show the linear trend of air temperature of +0.14 °C/100 year, trend of daily amount of precipitation of -0.30 mm/100 year. Curve of long-term salinity variation at Split-Gargano transect shows the increasing trend. This trend is evident also at the Kastela Bay coastal station.

Using some statistical method like Principal Component Analysis (PCA), time series of air temperature, sea temperature, pressure, precipitation and salinity were compared for coastal and open sea separately. Analysis allowed determination of main component important for variability of analysed climatic-oceanographic field on secular time scale.




Figure 1. Monthly mean time series filtered by 24M214 for station Trieste (northern Adriatic) for period 1840 - 1990

### REFERENCES

ZORE-ARMANDA, M., M. BONE, V. DADIC, M. MOROVIC, D. RATKOVIC, L. STOJANOSKI and I. VUKADIN, 1991. Hydrographic properties of the Adriatic sea in the period from 1971 through 1983. *Acta Adriat.* Vol. 32, 1: 1-547.

## MODELLING OF THE WIND DRIVEN CIRCULATION IN THE GULF OF LIONS

Stéphane DELEVILLE<sup>1</sup>, Claude MILLOT<sup>1</sup> and Laurent MORTIER<sup>2</sup>  
<sup>1</sup> Antenne C.O.M, B.P. 330, La Seyne-sur-mer, France  
<sup>2</sup> LODYC, Université de Paris 6, Paris, France

The Gulf of Lions is characterized all year long by the occurrence of gusty winds (the Mistral and the Tramontane) which, being constrained by the orography, blow from a constant direction (the North and the North-West resp.). As shown from both *in situ* and remotely sensed data (MILLOT, 1979) as well as from former numerical models (HUA, 1981) and specific analyses (CRÉPON *et al.*, 1984) dealing with summer conditions, the upwelling mainly occurs along straight 10-20 km segments of the coastline while the bottom topography doesn't play a major role. This semi-circular gulf is also under the influence of the Northern Current which continuously flows along the continental slope (i.e. along its diameter) and spreads over the shelf when the winds stop. Modelling such an area is thus relatively easy and useful to better understand the circulation in wider areas of upwelling such as North-West Africa, Peru,...

In this work, which is part of the EUROMODEL contribution to the MTP of MAST-2, we use the 3D primitive equations numerical model of LODYC with a cartesian grid and a quite realistic coastline. The domain, extending from Cap de Creux in the South-West to cap Sicié in the North-East (i.e. 300 km x 240 km) is such that a newtonian restoring on initial values of tracers and dynamics is used along the open boundaries to damp inertial waves. Due to the space scale of the phenomena and to the value of the internal radius of deformation (~9 km), the horizontal grid size is 3 km. The preliminary computations presented here are done with a constant depth of 100 m. For these particular studies, the model has a vertical resolution (15 levels) ranging from a few metres, allowing a correct representation of the mixed-layer and the thermocline (which has mean depth and thickness of 30 m and 10 m resp.) up to a few tens of metres at depth. These computations have been made with very simple wind conditions (direction of a Tramontane-like wind, constant speed of 9 m/s starting from rest), temperatures of 20°C in the mixed layer and 13°C at depth, and without any general circulation.

After three inertial periods (about 3 x 17.5 h), which is a characteristic time scale for the summer gusts of wind, the distribution of the surface temperature is very similar to the commonly observed one. Upwellings are obviously located along the coast on the left of the wind, and mainly along the straight coastal segments, thus showing a marked discontinuity due to the coastline geometry. The intensity is also different from one upwelling to another, with the lowest temperatures encountered near the coast ranging from 4°C to 5°C below the initial surface temperature. All these features are in good agreement with *in situ* and remotely sensed observations.

The circulation in the surface layer forms an eastward coastal jet of ~50 cm/s along the northern coast. Capes located on the windward side of the straight coastal segments induce a very local intensification of the jet. In the central part of the gulf, the wind driven circulation is almost on the right hand side of the wind. At the bottom, the coastal circulation is characterized by a westward jet along the northern coast and a northward one along the western coast, each being of a few cm/s. Due to the very large theoretical depth, the currents in most of the gulf are non significant. Most of these features are in agreement with the available observations.

After this study of the effect of a Tramontane-like wind, some experiments will be run with a Mistral-like wind and then with a combination of both. We want to analyse the effects of the wind strength by increasing its speed (from ~9 m/s to ~15 m/s). Some experiments will be done with a time depending wind to see the role of the succession of the gust of winds in the formation of upwellings and this study will end with actual wind fields, provided by meteorological numerical models such as PERIDOT.

This study is a part of a program of modelling the seasonal and meso-scale variations of the circulation in the Gulf of Lions: we want to study the interactions with the general mediterranean circulation (HERBAUT *et al.*, 1994).

### REFERENCES

CRÉPON M., C. RICHEZ and M. CHARTIER, 1984. Effects of coastline geometry on upwellings. *Journal of Physical Oceanography*, Vol. 14, 8: 1365-1382.  
HERBAUT C., L. MORTIER, M. CRÉPON, 1994. A sensitivity study of the general circulation of the Western Mediterranean Sea. Part 1: The response to density forcing through straits. Submitted to *Journal of Physical Oceanography*.  
HUA B.L., 1981. Modélisation numérique d'upwellings côtiers à l'aide d'une méthode d'éléments finis. Application au Golfe du Lion. Thèse d'état Paris 6, 213 pp.  
MILLOT C., 1979. Wind induced upwelling in the Gulf of Lions. *Oceanol. Acta* 2, 3: 261-274