

CETACEAN SIGHTINGS IN THE ALBORAN SEA. JULY 1993

FRANCO I.¹, MAS J.² and RODRIGUEZ C.²

¹ IEO. Centro costero de Fuengirola. Ap. C. 285. 29640-Fuengirola, Málaga, España
² IEO. Centro Costero de Murcia. Ap. C. 22. 30740 S. P. del Pinatar, Murcia, España

The Mediterranean cetacean populations have in recent years been surveyed by several authors. These surveys have mainly taken place in the sea water around Italy: Ligurian sea (GANNIER & GANNIER, 1990; FABRI & LAURIANO, 1990), Tyrrhenian sea (CONSIGLIO *et al.*, 1990), Corsica, Sardinia and Sicily (NOTARBARTOLO *et al.*, 1990), Adriatic sea (SERMAN & SERMAN, 1990), in the Ionic sea (POLITI *et al.*, 1990) and in the northern western Mediterranean (AGUILAR, 1990; AGUILAR *et al.*, 1992; FORCADA *et al.*, 1990) and only a few studies have taken place in the Alborán sea (FORCADA *et al.*, 1990; AGUILAR *et al.*, 1992). From 8 to 27 July 1993, a cetacean survey cruise took place between 6°12'00"W and 2°0'00"W in the Alborán Sea and Gibraltar Strait waters. The observation platform was placed at 5 m. above sea level. A total of 127 hours of observations were made on board of the R/V Francisco de Paula Navarro during the IEO ICTIO ALBORAN 0793 cruise, covering a distance of 1288 n. miles (Fig. 1) with an area of 13.213 n. mi². Line transect sampling methods (BURNHAM *et al.*, 1984) were used to analyze the data and to calculate the estimation of abundance for the whole area. Along with the date and time sightings, position, species, number of individuals, depth, distance to the coast, and distance to the sighting data of sea conditions, temperature, atmospheric conditions and visibility were also recorded together data of the cetacean behavior, school type, speed and course, swimming behavior, attraction to the vessel, the birds and to the other fauna. The total number of sightings was 62. The species encountered were *Delphinus delphis* (31%), *Globicephala melas* (26%), *Stenella coeruleoalba* (23%), *Tursiops truncatus* (18%), *Physeter macrocephalus* (1%) and *Grampus griseus* (1%). The LDS (density of sightings) was calculated giving a value of 0,048 schools/n.mile. Estimates for grouping index, distance to the coast, behavior, temperature ranges, etc., for the different species are given in table I.




Fig. 1. Area surveyed and situation of the sightings.

REFERENCES

AGUILAR A., 1990. Calvin in the striped dolphin from the western Mediterranean sea. pp. 67-68. In European Research on Cetaceans. - 4. Proc. 4 Th. Ann. E. C. S. Conf. AGUILAR R., PASTOR X. & FORCADA J. 1992. Results of the Greenpeace Cetacean Survey Cruises during the western Mediterranean striped dolphin epizootic. BURNHAM K.F., ANDERSON D. R. & LAAKE J. L. 1984. Estimation of density from line transect sampling of biological populations. Wildlife monogr. 72: 1-202. CONSIGLIO C., MARINI L., ANGRADI A. & SANNA A., 1990. A medium-term sighting scheme on cetacean in the central Tyrrhenian sea. pp. 44-45. In European Research on Cetaceans. *op. cit.* FABRI F. & LAURIANO G., 1990. Greenpeace report on two year research in the Ligurian sea. pp. 69-71. In European Research on Cetaceans. *op. cit.* FORCADA J., AGUILAR A., HAMMOND P., PASTOR X. & AGUILAR, R., 1992. Striped dolphin abundance in the Western Mediterranean Basin after 1990 epizootic. The Mediterranean Striped Dolphin Die-off. Proc. of the Mediterranean striped dolphin mortality International Workshop, Palma de Mallorca. GANNIER A. & GANNIER G., 1990. Cetacean sighting in the Mediterranean sea. Second Report. pp. 39. In European Research on Cetaceans. *op. cit.* MARINI L., CONSIGLIO C., ANGRADI A., SANNA A. & VALENTINI T., 1990. Cetacean sightings programme in the central Tyrrhenian sea. pp. 66-67. In European Research on Cetaceans. *op. cit.* NOTARBARTOLO G., VENTURINO M.C., ZANARDELLI M., BEARZI G., BORSANI F., CAVALLONI B., CUSSINO E., JAMODA M. & ATROLDI S., 1990. Distribution and relative abundance of cetaceans in the central Mediterranean sea. pp. 41-43. In European Research on Cetaceans. *op. cit.* POLITI E., BEARZI M., NOTARBARTOLO DE SCIARA G., CUSSINO E. & GNONE G., 1990. Distribution and frequency of cetaceans in the waters adjacent the greek ionian islands. pp. 75-78. In European Research on Cetaceans. *op. cit.* SERMAN D. & SERMAN A., 1990. Marine mammal conservation status and research in the eastern Adriatic sea. pp. 54-55. In European Research on Cetaceans. *op. cit.*

Table I. Summary of all data obtained in the sightings.

	D. del.	G. mel.	S. coe.	T. tru.	P. mac.	G. gri
n° sightings	19	16	14	11	1	1
n° individuals	236	118	129	57	1	1
LDS	0,014	0,012	0,01	0,008	0,0008	0,0008
Compact groups	1	6	1	1	-	-
Sparse groups	18	10	13	10	-	-
Isolated individuals	0	0	0	0	1	1
Average n° individ./group	12,42	7,37	9,21	5,18	1	1
Attraction to the vessel	12	6	3	6	-	-
Move away from the vessel	0	1	0	0	-	-
Indifferent to the vessel	7	9	11	5	1	1
Stationary swimming	2	7	5	1	-	-
Slow swimming	13	12	13	8	1	1
Fast swimming	6	0	0	3	-	-
Showing the back	18	16	13	10	1	1
Showing the caudal fin	4	0	6	2	-	-
Jumping	15	0	7	1	-	-
Maximum depth	1300	1500	1400	1150	700	1300
Minimum depth	80	375	75	30	700	1300
Average depth	685	869	727	407	700	1300
Max. distance to the coast	2,2	10,1	3,8	1,0	23,8	34
Min. distance to the coast	39,8	39,6	42,0	32,0	23,8	34
Ave. distance to the coast	15,9	24,7	18,8	13,7	23,8	34
Max. water temperature	21,3	23,9	23,6	23,2	20,8	21,8
Min. water temperature	18,6	21,4	21,3	20,8	20,8	21,8
Ave. water temperature	20,2	22,4	22,4	21,8	20,8	21,8

DAILY EGG PRODUCTION SPAWNING BIOMASS OF THE NORTH-WESTERN MEDITERRANEAN ANCHOVY DURING 1993 (CATALAN SEA, GULF OF LIONS AND LIGURIAN-N TYRRHENIAN SEAS)

A. GARCÍA¹, I. PALOMERA², J.P. RUBÍN¹, N. PÉREZ¹, O. GIOVANARDI³ and P. RUBIÉS²

¹ IEO. Centro Oceanográfico de Málaga, P.O. Box 285. Fuengirola, Málaga, Spain
² Instituto de Ciencias del Mar (ICM), Paseo Joan de Borbó s/n, 08039 Barcelona, Spain
³ Ist. Centrale Per La Ricerca Scientifica e Tecnologica applicata al Mare, Roma, Italy

Research carried out by HUNTER AND GOLDBERG (1980) on the reproductive aspects of the northern anchovy off California *Engraulis mordax*, showed that many small pelagic species, mainly clupeiforms were batch spawners. Subsequently, an ichthyoplankton based method (LASKER, 1985), the Daily Egg Production Method, was developed that allowed the spawning biomass estimate of Clupeoids. The first applications of DEPM in European waters were carried out in the coasts of the Atlantic Iberian peninsula on the Iberian sardine, *Sardina pilchardus* (GARCÍA *et al.*, 1992) during 1988, and on the Bay of Biscay anchovy, *Engraulis encrasicolus* (SANTIAGO and SANZ, 1992). The first Mediterranean DEPM survey was carried out by PALOMERA and PERTIERRA (1993) on the anchovy from the Catalan coasts. Within the framework of an EU financed FAR project, the northwestern Mediterranean anchovy biomass was estimated through DEPM (MPH-MED-93) aboard the R/V García del Cid during July, 1993. This survey was combined spatially and temporally with the echo-acoustic survey, PELMED-93 on board the R/V Thalassa, with the purpose of integrating the anchovy echo-integration biomass estimates and allowing to estimate DEPM parameters related to the adult stock. The DEPM sampling and data treatment methodology in relation to the egg and adults survey are described in GARCÍA (1994). In summary, the basic scheme of egg sampling stations was based on a 5 by 5 nautical mile track, with transects perpendicular to the coastline. A total of 602 CalVET net (150 µ mesh) vertical tows of 100 m depth were done, representing a coverage of 59,981 km² of sea surface. Catalan Sea accounted for 292 plankton hauls, whereas the Gulf of Lions and Ligurian Sea accounted for 138 and 172, respectively.

Adult anchovies were sampled in 34 positive anchovy hauls (13 in Catalan waters, 13 in the Gulf of Lions and 8 in the Ligurian-N Tyrrhenian) with an epipelagic trawl. The period of trawls ranged from 7:30 A.M. to 22:30 P.M. (GMT). 1,034 ovaries were collected, where 210 corresponded to hydrated females. The model (LASKER, 1985) is based on the following equation, where B = spawning biomass in metric tons, Po = daily egg production (number of eggs per sampling unit, 0.05 m²), W = average weight of mature females (grams), R = sex ratio (fraction mature of females by weight), F = batch fecundity (mean number of eggs per mature female per spawning), A = total survey area (in 0.05 m² sampling units), S = fraction of mature females spawning per day. Plankton stations are post-stratified by location (negative and positive strata) with the purpose of decreasing variance and by geographic criteria based on the spawning area distribution (GARCÍA, 1994). Eggs were staged according to their embryonic degree of development and subsequently aged, taking into account an specific temperature-dependent egg development model, through the program STAGEAGE. Egg production, Po, is estimated by fitting the exponential mortality function using a weighted nonlinear least squares regression to the data egg file. This model was fit to the data from stratum 1 for each of the regions. In consequence, each region has an estimate of Po1 (intercept) and a corresponding egg mortality, z (slope). The final stratified estimate of Po by regions was calculated as the weighted average of the two strata. In reference to the adult parameters estimates, W, F, S and R, mean and variance were estimated following PICQUELLE and STAUFFER's (1985) weighing procedure, calculating weighted averages albeit the number of sampled individuals are not equal in each of the tows. W, the mean weight of mature females per trawl was adjusted for those females which were in the hydrated condition, through the following regression, $W = -0,3261 + 1,1012W^*$ (ovary-free weight of non-hydrated females). F, batch fecundity, is estimated by regressing batch fecundity on ovary-free weight of hydrated females, without post-ovulatory follicles. A total of 83 hydrated females were obtained. The resulting linear regression is: $y = 1848,3 + 499,57 x$ R = (0,92). Weights of the females used ranged from 3.6 to 30.6 g, while sizes ranged from 9.1 to 17.5 cm. S, fraction of mature females spawning per day, is determined by the histological analysis of post-ovulatory follicles. These were assigned ages according to the following: Day-0 PO = 0-7 hours; Day-1 PO = 8-31 hours; Day-2 PO = > 31 hours. Day-1 PO has been used in the estimate of S, since Day-0 PO are oversampled during the daily spawning period in anchovy (ALHEIT, 1985). R, sex ratio, is calculated as the fraction in weight of mature females. Based on the geographical distribution of the anchovy population and the spawning grounds distribution, the defined stratified regions were: Catalan Sea and Gulf of Lions, which have common geographical, hydrographical and environmental characteristics, as opposed to the Ligurian-N Tyrrhenian Sea, which presents a

DEPM Parameter	Catalan Sea-Gulf of Lions	Ligurian-N Tyrrhenian Sea
Po1	0.215	4.350
z	0.145	0.223
W	0.003	1.561
F	0.136	0.290
Po	0.342	0.716
S	0.295	0.225
Act. (0.05 m ²)	6,911 x 10 ¹¹	3,085 x 10 ¹¹
Act. (km ²)	44,567	15,423
W (g)	14.31	14.17
F	0.172	0.058
S	0.11	0.16
R	0.31	0.30
z	0.12	0.11
R	0.85	0.80
S	0.25	0.25
Biomass (MT)	30,565	12,129
z	0.23	0.30
Acoustic Biomass (MT)	32,831	6,459

distributional barrier along the Provençal coasts, probably due to the narrow shelf limits. The results of all the DEPM parameters estimate and the final anchovy spawning biomass estimate are in following table. When comparing by regions the different parameters estimate, no great differences are observed between the adult parameters. However, in comparison to the Bay of Biscay anchovy, it should be remarked that the average female weight is much lower and consequently, batch fecundity (approximately 30 g and 15 000 eggs/batch, respectively). Nevertheless, spawning fraction is within the same range (0.30). Daily egg production over the surveyed regions are in the same order of magnitude, but with high mortality (z), specially in the Ligurian-N Tyrrhenian Sea, in comparison to Bay of Biscay anchovy estimates (SANTIAGO and SANZ, 1992). Higher Mediterranean temperatures account for the faster egg development rates that eventually result in shorter egg durations. The temporal and spatial simultaneous coverage of both surveys have contributed to the similarities in the biomass estimates, except the case of the Italian waters in which daily egg production probably is over-estimated.

REFERENCES

ALHEIT, J., 1985. Technical Rep. NMFS 36:59-62.
 GARCÍA, A. (Ed.), 1994. Final Report Study Contract FAR Project MA 3.730. U.E. DG XIV
 GARCÍA, A., N. PÉREZ, N.C.H. LO, A. LAGO DE LANZOS and A. SOLA, 1992. *Bol. Inst. Esp. Oceanogr.*, V. 8, N° 1: 123-138.
 HUNTER, J.R. and S.R. GOLDBERG, 1980. *Fish. Bull. U.S.* 77: 641-652.
 LASKER, R. (Editor), 1985. *NOAA Technical Rep. NMFS* 36: 99 p.
 PALOMERA, I. and PERTIERRA, J.P., 1993. *Sci. Mar.*, 57(2-3): 243-251.
 PICQUELLE, S.J. and G. STAUFFER, 1985. *NOAA Technical Rep. NMFS* 36:7-16.
 SANTIAGO, J. and A. SANZ, 1992. *Bol. Inst. Esp. Oceanogr.*, V. 8, N° 1: 225-230.