

MONITORING FOR PROTECTION OF THE MARINE ENVIRONMENT USING LANDSAT-TM DATA

A.P. CRACKNELL¹, K. ABDULLAH¹,
J. N. HATZOPOULOS², M. KARYDIS², D. GAZIS²

¹ Depart. of Applied Physics Electronic & Manufacturing Engin., Univ. of Dundee, Scotland
² Department of Environmental Studies, University of the Aegean, Mytilene, Greece

This paper presents results from a joint project between the University of Dundee and the University of the Aegean. The project deals with the monitoring of the quality of the sea water environment using *in situ* measurements and satellite image data. The study was carried out using Landsat-TM data in Mytilene harbour and the surrounding water areas. *In situ* water samples were collected and analysed at the University of the Aegean during the year 1992 for the dates : 5 March, 9 June, 11 July and 28 August. Water samples were collected with a Van Dorn sampler from the depth 0-1m. The water samples were used for chlorophyll and suspended matter determinations. Temperature and salinity was measured by a CTD instrument Model YSI 6000. Light transparency was recorded using a secchi disk and a KALSHICO digital underwater irradiameter model 268 WA 305. Samples for chlorophyll determinations were filtered immediately through millipore filters (pore size 0.45 um) after collection and the pigments of the filter were extracted overnight in 90% acetone. The samples were then centrifuged (3000 g for 30 min) and the chlorophyll absorption was measured in a double beam spectrometer Model Varian DMS-80. The detailed procedure is described by elsewhere (UNESCO-SCOR, 1966).

Suspended matter was measured gravimetrically according to the procedure described by STRICKLAND & PARSONS (1972). Estimates of suspended matter were also

carried out by turbidity measurements using a Hach turbidity meter. Four Landsat-TM cloudless mini scenes (50x50 km coverage) were purchased corresponding to the *in situ* sampling dates. The position of the sampling boat was determined by photogrammetry and later by GPS. The image processing was performed at the University of Dundee (CRACKNELL *et al.*, 1994) in the following steps: (a) image rectification to a UTM 1:25000 scale map, (b) transformation from the sample location to image pixel scan lines, (c) the pixel values for bands 1, 2 and 3 were extracted for a 3 pixel by 3 pixel area corresponding to the sites of the *in situ* measurements and then they were converted to (i) radiance values and (ii) exo atmospheric reflectance values, and, the atmospheric correction was performed to the data using the darkest pixel method, (d) transformations such as the principal component, characteristic vector (using reflectance data) and chromaticity (using radiance data) were performed on the data sets, (e) multiple regression analyses were performed with dependent variables chlorophyll and suspended matter concentrations and their reflectance values. Two algorithms were then derived given by the Equations :

$$y = a + b_1r_1 + b_2r_2 + b_3r_3 \quad (1)$$

$$y = a + \sum_{i=1}^3 b_i r_i + b_4 r_1 r_2 + b_5 r_1 r_3 + b_6 r_1 r_3 \quad (2)$$

Where y is the concentration of chlorophyll or suspended matter, r_i is the reflectance value for band i and the coefficients a, b_i ($i=1$ to 6) are selected empirically and determined by the regression. Water quality maps were then generated by applying Equation (2) to all sets of data except June where Equation (1) was used. The distribution patterns of chlorophyll show that in March (Figure 1) a higher concentration occurred in the vicinity of Mytilene harbour and at some locations along the coast to the north and also in the coastal waters in the north eastern part of Kolpos Geras. Figure 2 shows the chlorophyll distribution for 11 July. Separate processing was applied to TM images of channel 6 to produce sea surface temperature (SST) maps which are shown in Figure 3 for the 5 March data set and Figure 4 for the 11 July data set. The algorithm used had the following form for all data sets (T_{sat} is the TM channel 6 value) :

$$SST = 11.1620 - 11.3132 \log(T_{sat}) \quad (3)$$

This equation was produced by regression with correlation coefficient of 0.98 and an RMS error of $\pm 0.8^\circ\text{C}$

REFERENCES

- CRACKNELL A.P. K. ABDULLAH, J.N. HATZOPOULOS, M. KARYDIS and D. GAZIS, 1994 : Monitoring for Protection of the Marine Environment Using Landsat TM and AVHRR Data, Final Report to the British Council of Athens, Greece.
STRICKLAND J.D.H. & T.R. PARSONS, 1972 : A Practical Handbook of Sea Water Analysis. *Fish. Res. Bd. of Canada*, 167 (2nd Ed.).
UNESCO-SCOR, 1966 : Monographs on Oceanographic Methodology, I. Determination of Photosynthetic Pigments in Sea Water, UNESCO, Paris, pp. 69.




Figure 1. Map of Chlorophyll Distribution (mg m^{-3}) on 5 March 1992. Colour code: blue <0.1 ; green $0.1-0.4$; yellow $0.4-0.7$; orange $0.7-0.1$; and red >1.1 .



Figure 2. Temperature Distribution ($^\circ\text{C}$) on 5 March 1992. Colour code: blue $12-13$; green $13-14$; yellow $14-15$ and red >15 .