

TEXTURAL AND COMPOSITIONAL CHARACTERISTICS OF SUSPENDED MATTER IN THE LIGNANO BASIN (MARANO LAGOON, NORTHERN ADRIATIC SEA)

G.P. FANZUTTI, E. COLIZZA, F. FINOCCHIARO, R. MELIS and R. PIANI

Istituto di Geologia e Paleontologia, Università di Trieste, Italia

The aim of this study is the evaluation of the mean characteristics of the suspended matter of the Lignano basin (surface: 40 km²; mean depth: 0.8 m), in the Marano Lagoon (Northern Adriatic Sea). Nine stations were sampled four times during 1991-1992 (October, February, April and July) along the channels of the basin, at lagoon inlet and at the Stella River mouth, which is the main spring-river, tributary of the lagoon (annual average discharge: 34 m³/s). Surface and bottom samples were taken with a Niskin bottle, twice a day, during flood and ebb in spring tides without wind conditions. Temperature, salinity (CTD probe) and current speed measurements (NBA currentmeter) were also collected.

The main parameters like total suspended matter (TSM), total carbonates and particulate organic carbon (POC), C/N ratio, 5th percentile, modal and mean diameter are here presented. It was established their most frequent values so as their exceptional ones, showing the differences among the various lagoon environments: the basin, the inlet and the mouth of the Stella River.

For almost all the parameters a systematic difference between the surface and the bottom was observed: concentration, mean diameter and carbonate percent values observed at surface are generally lower than those at the bottom. Likewise at the lagoon inlet (FANZUTTI *et al.*, 1992), this difference could be explained by the selective sedimentation within the water column of lagoon channels.

By observing the TSM frequency distribution, it can be established that the most frequent values are included in the band between 4 and 10 mg/dm³ for the lagoon basin, those between 12 and 30 mg/dm³ become less frequent, and those above 30 mg/dm³ should be considered unusual. The last ones can be found mostly in surface during February and along the water column during July.

At each sampling, it was observed that the mean concentration values in the inlet are usually lower than those measured in the lagoon basin, whereas in the spring-river water values are similar to those of the lagoon. The only exception appears in October when the Stella River discharge was higher than the annual average and when at the inlet the resuspension phenomena due to the sea state are verified.

The frequency distribution of the POC in the lagoon basin shows a modal interval between 250 and 1000 µg/dm³. Values lower than 250 µg/dm³ are not found. The frequency of values higher than 1000 µg/dm³ tend to decrease roughly, and likewise it is observed for the TSM, very high values at surface during February and along the water column during July are observed. At each sampling the lagoon inlet showed values equal or lower than those observed in the lagoon, while Stella River mouth (in normally discharge conditions) presents organic carbon values lower than those of the lagoon. In the lagoon, modal C/N ratio values range between 8 and 10.

The mean percent values of detrital carbonates in the basin are between 31.6% and 39.5%, while the range in the inlet is widely (29.5% in February and 42.5% in July). In the mouth of the Stella River the carbonate contents are slightly superior to those of the lagoon basin.

In the lagoon basin, the frequency distribution of grain-size parameters has allowed to establish that the most frequent values of the mean diameter are between 8 and 12 µm. As far as the modal diameter is concerned, values range from 10 to 12 µm, whereas for the 5th coarse percentile between 24 and 26 µm. Among all these parameters, the mean diameter is the one that permits to differentiate the various environments. During ebb tide, the mean diameter in the inlet are higher with respect to the basin one, while during the flood the opposite phenomenon is observed.

The particulate matter brought from the Stella River is coarser (10 to 14 µm) than the lagoon one. This phenomenon is more evident in October sampling when the annual mean discharge was exceeded. Whether for the characteristics of the suspended matter or for the salinity distribution, the diffusion of the "plume" within the lagoon seems to be limited to a distance of about 2 km from the mouth (BELLI *et al.*, 1994).

BRAMBATI *et al.* (1990) recognized that a resuspension forced by the wind is one of the factors that mostly change the patterns of the TSM concentration in the Lignano basin. Since the data here presented are not influenced by wind resuspension, it was possible to indicate normal values during spring tides, while the exceptional values could be attributed to others phenomena below mentioned.

Anomalous values of POC found in February should be attributed to organic debris, as confirmed by the high values of C/N ratio and the low values of chlorophyll (FONDA UMANI, personal communication). The accumulation of organic debris within the basin could be due to the flood, typical of the late autumn season and to the degradation of the macro-algae that colonize the tidal flat. Its drastical removal is caused by the maximum (equinoctial) February tide excursions when wide zones of tidal flat emerge. The high concentrations of TSM found in July should be related to the resuspension caused by an intensive traffic of boats, which highly increases during the summer time in the resort area of Lignano.

This work is a step of a research that involves also ANPA-DISP (Rome) and Laboratorio di Biologia Marina (Trieste). It is supported by ENEA-DISP and MURST 60% "Ambienti Umidi" grants.

REFERENCES

- BRAMBATI A., FANZUTTI G.P. and FINOCCHIARO F., 1990. Effetti della risospensione indotta da vento sulle concentrazioni e dimensioni del particolato nel bacino di Lignano (Laguna di Marano-Adriatico settentrionale). *Atti 8° Congr. AIOL*: 191-212.
- FANZUTTI G.P., FINOCCHIARO F. and PIANI R., 1992. A comparison among some suspended matter characteristics in two tidal inlets of the Marano Lagoon (Northern Adriatic Sea). *CIESM, Rapp. Comm. Int. Mer Médit.*, 33, 400.
- BELLI M., COLIZZA E., FANZUTTI G.P., FINOCCHIARO F., MELIS R., PIANI R. and SANSONE U., 1994. The role of a spring river as a source of Radiocaesium in a lagoonal environment: the case of the Stella river (Marano Lagoon, Northern Adriatic Sea). *Int. Seminar on Freshwater and Estuarine Radioecology*, Lisboa, 21-25 March 1994.