

DIATOMS IN HYPERSALINE SOLAR SALTERN PONDS (EBRO DELTA, SPAIN)

MERINO V.¹, CLAVERO E.², GRIMALT J.O.² & HERNÁNDEZ-MARINÉ M.¹

¹ Lab. of Botany. Fac. of Pharmacy. Univ. of Barcelona. Av. Joan XXIII, s/n.

² Dept. of Environmental Chemistry (CID-CSIC). C/Jordi Girona, 18.
08034 Barcelona. Spain.

Benthic diatoms constitute an important component of the microbial mats developed in the lower range of salinities in hypersaline environments, although few studies on diatom communities in such environments have been published to date (EHRlich & DOR, 1985, NOËL, 1984). This paper presents the diatoms that thrive in the solar saltern ponds "La Trinitat" and their distribution as function of salinity.

In these salt fields (Ebro delta, 40°35'N, 0°40'E, South Catalonia, Spain) halite is obtained by pumping sea water to an array of evaporation ponds. The water flows through this system, and evaporates and increases in salinity through successive ponds. The salt concentration within each pond changes only slightly owing to human control, hence communities of each lagoon develop in a high but constant salinities. After the halite is harvested, from August through November, the pools are once again flooded with sea water.

Diatoms appear in the first three ponds of the circuit, with salinity ranging from that of sea water to 120‰. In these ponds (called Deposits 0, 1 and 2), sedimentation of sand and organic matter and carbonate precipitation occurs (CLAVERO *et al.*, this volume). Samples for optical, SEM and TEM examination and measurements of physical and chemical parameters were obtained monthly in 1994. When possible, depending on the presence and abundance of photosynthetic organisms, the sediment was sampled by taking cores. These cores were sliced in roughly 1 mm-thick sections, corresponding to the sediment layers.

In the lowest salinity pond, Deposit 0 (mean salinity 48‰), either the floor was nude or the organisms did not produce layers. Under the shallow water, *Ruppia maritima*, *Chaetomorpha* sp. and *Cladophora* sp. developed in spots. Some diatom species, e.g., *Striatella unipunctata* (Lyngbye) Agardh, *Achnantes brevipes* Agardh, *Cocconeis placentula* var *euglypta* (Ehr.) Cleve, appeared in high quantities among the macrophytes or epiphytizing them. Surrounding the pond, in spring, there is a temporarily flooded ring with a water covering of only few mm, in which a distinct laminated mat of filamentous cyanophytes develops with a golden-brown upper layer built by diatoms. This community was dominated by species of the genera *Amphora*, *Navicula* and *Nitzschia*: *Nitzschia lembiformis* Meister, *Nitzschia sigma* (Kütz.) W. Smith and *Nitzschia vidovichii* (Grun.) Peragallo. As summer went on the marginal ring dried up with a leathery appearance.

In Deposit 1 (mean salinity 64‰) the most abundant diatom was *Pleurosigma elongatum* W. Smith, which appeared abundantly on the surface sediment, associated with filaments of cyanophyta. It also appeared on sporadically wet sediments and forming floating lumps together with *Lyngbya aestuarii* Liebm. Other diatoms that also appeared in abundance were *Scolioleptura tumida* (Brev.) Rabenhorst and some species of *Navicula* and *Amphora*: *A. coffeaeformis* (Agardh) Kützing, *A. acutiuscula* Hustedt, *A. hyalina* Kützing. *Cocconeis placentula* var *euglypta*, *Pleurosigma elongatum* and *Nitzschia sigma* were also observed associated with *Cladophora* sp. In summer, the partially flooded borders developed a green and white mass of *Beggiatoa* sp. and *L. aestuarii* mixed with living *Surirella striatula* Turpin and *Pleurosigma elongatum*.

In Deposit 2 (mean salinity 97‰) the number of species and individuals was smaller. The most abundant was *Nitzschia lembiformis* (which also appeared in Deposit 1, but less frequently). *Surirella striatula*, *Amphora coffeaeformis*, *Nitzschia frustulum* (Kützing) Grunow and *Nitzschia sigma* were also well represented.

Heaters (subsequent ponds with salinities over 120‰) were devoid of diatoms, except for empty valves and girdle bands of *Nitzschia* sp., probably allochthonous.

Some of the taxa found in the least saline pond (Deposit 0) are known from the nearby marine waters, e.g., *Striatella unipunctata*, *Surirella fastuosa* (Ehr.) Kützing, and *Achnantes brevipes*, and were not observed in ponds of higher salinities. Others, like *Nitzschia vidovichii*, *Pleurosigma elongatum* and *Gyrosigma spencerii* (Quek.) Grif. et Henfr., occurred only in waters with 44-80‰. *S. Nitzschia lembiformis* was found only with salinities from 53‰ to 150‰ predominantly in the hypersaline pond Deposit 2 (85-105‰ S). The salinity range gives this species an "hyperhalobius" character (EHRlich & DOR, 1985). Most of the diatoms recorded were euryhaline, but their halotolerance varied. A general decrease of diatom diversity in increasing salinity was found. Some species disappeared at lower salinities, while several species were present throughout the entire area at salinities of 44-115‰, e.g., *Amphora angusta* (Gregory) Cleve, *A. coffeaeformis*, *Nitzschia frustulum* and *Nitzschia sigma*, in accordance with their classification as euryhaline forms (EHRlich & DOR, 1985, NOËL, 1984).

A remarkable fact is that no frustules of central diatoms were found in sediments, which points out that in these shallow saline waters central diatoms did not develop, which agrees with previous results (NOËL, 1984).

REFERENCES

- EHRlich A. & DOR I., 1985.- Photosynthetic microorganisms of the Gavish Sabkha. In: Friedman & Krumbein (eds.) Hypersaline Ecosystems. The Gavish Sabkha: 296-321.
CLAVERO, E., MERINO, V., GRIMALT, J.O. & HERNÁNDEZ-MARINÉ, M., 1995.- Cyanophyta in hypersaline solar saltern ponds (Ebro Delta, Spain). *Rapp. Comm. int. Mer Médit.*, this volume.
NOËL D., 1984.- Les diatomées des saumures et des sédiments de surface du Salin de Bras del Port (Santa Pola, province d'Alicante, Espagne). *Revista d'Investigacions Geològiques*, 38/39 : 79-107.