

EFFECT OF SOME ENVIRONMENTAL AND PHYSIOLOGICAL FACTORS ON REPRODUCTIVE PARAMETERS OF *MUGIL CAPITO* DURING THE BREEDING SEASON

ZAKI M.I., EL-GREISY A.A., EL-GHARABAWY M.M. and BAGHDADI H.H.
National Institute of Oceanography and Fisheries, Alexandria, Egypt

The effect of some environmental and physiological factors on reproductive parameters of *Mugil capito* in Egyptian fresh water fish farm has been studied. The present investigation revealed that the increase in photoperiod (18L+6D) as well as continuous illumination (24L) and continuous darkness (24D), showed an obvious decrease in the gonadosomatic index (GSI) compared to that of the control (6,5L+17,5D). This result is in affirmative agreement with that observed by LAM and SOH (1978) on *Siganus canaliculatus*.

The present study also revealed that the increase in temperature showed an obvious decrease in the GSI of both sexes of *Mugil capito* while the decrease in the temperature showed a decrease in the GSI of the female and a slight increase in the GSI of the male.

The GSI of the females increased within 1 week of 1500 IU injection. After 10 weeks, the gonads were resorbed and the GSI greatly decreased. There is no sharp difference in the GSI of the males after the injection.

The egg diameter of *Mugil capito* is affected greatly by the change in the environmental factors. The present study showed that the egg diameter is inversely proportional to the photoperiod.

From the relation between diameter and temperature it is observed that the increase or decrease in temperature is accompanied by a decrease in the values of egg diameter. In both cases, the frequency distribution was shifted to the lower values compared to that of the control. This result with that observed by TAMARU *et al.*, (1991) who studied the egg diameter of *Mugil capito*.

Concerning the effect of HCG injection on the egg diameter, it was observed that the frequency distribution of egg diameter was shifted to the longer diameter after 1 week of the injection. At the end of the experiment (10 weeks) it was shifted to the shorter one. The same increase in oocyte diameter was observed in the stimulated females of black porgy, *Acanthopragus schlegeli* due to the injection of HCG (CHANG *et al.*, 1991).

Milt volume in the group that received the highest dose (2 000 IU HCG) was greater than in the other group that received the lower dose (1 500 IU HCG). The present study also revealed that the sperm concentration in the seminal fluid decreased with time and with the number of times for the fish to give milt.

REFERENCES

- CHANG C.F., YUEH W.S. and LEE M.F. 1991. Effects of LHRH-A and HCG on the steroid profiles of bisexual and mature male and female protandrous black porgy, *Acanthopragus schlegeli*. *Aquaculture*, 92 : 83-92.
LAM Y.H. and SOH C.L. 1975. Effect of photoperiod on gonadal maturation in the rabbit fish *Siagnus canaliculatus*, park 1797. *Aquaculture*, 5 : 407-410.
TAMARU C.S., KELLY C.D., LEE C.S., AIDA H. HANYU I. and GOETZ F. 1991. Steroid profiles during maturation and induced spawning of the striped mullet, *Mugil cephalus* L.. *Aquaculture*, 95 : 149-168.