


BOTTOM SEDIMENT DISTRIBUTION IN THE GULF OF CÁDIZ SLOPE INFLUENCED BY THE MEDITERRANEAN UNDERCURRENT

RODERO J.¹ and MALDONADO^{1,2} A.

¹ Instituto Andaluz de Geología Mediterránea, CSIC. C/Fuentenueva s/n 18002.

² Depart. de Geodinámica, Universidad de Granada. C/Fuentenueva s/n 18002.
Granada, Spain

As a result of the opening of the strait of Gibraltar at the end of the Miocene, the connections between the desiccated Mediterranean sea and the Atlantic ocean were established with circulation patterns probably similar to the present (MALDONADO, 1992). The interaction between different water masses resulted in a complex current system in the Gulf of Cádiz continental margin, which influenced sediment distribution and depositional patterns (NELSON *et al.*, 1992). One of the main controlling factors of the sediment distribution in the Gulf of Cádiz is the interplay of several water masses at different depths in response to the circulation patterns of the flows from the Mediterranean and the Atlantic water masses (HEEZEN and JOHNSON, 1979). The flow from the Mediterranean is known as the Mediterranean undercurrent, which is characterised by high salinity, resulting in a more dense water mass than the colder Atlantic waters. As it leaves the strait of Gibraltar, the Mediterranean undercurrent flows between 200 and 1800 m water in a WNW to NW direction along the continental margin of the Gulf of Cádiz, influencing particularly sediment distribution along the shelf brake and upper slope. The predominant Atlantic water flow, in contrast, follows the continental shelf in a ESE to SE direction to enter the Mediterranean sea through the strait of Gibraltar.



The Mediterranean undercurrent decreases in speed as it moves NW, away from the strait of Gibraltar. The speed ranges between maximum velocities of 250 cm/s (KENYON and BELDERSON, 1973) to 181 cm/s (AMBAR and HOWE, 1979) near the strait, to 10-20 cm/s in the central sector of the Gulf of Cádiz. This decrease in energy results in a distinct gradation of sediment types in the margin. Four main areas have been identified in the continental margin on the basis of the grain-size distribution:

Area 1. This area occupies the southernmost sector of the Gulf of Cádiz. It is characterised by bioclastic sand and gravel, developed as a lag deposit due to the erosion and transport of the finer, terrigenous fraction by the strong Mediterranean undercurrent. There are also locally outcrops of the basement, without sediment cover due to the strong bottom current. These sediments deposits develop several types of bedforms (NELSON *et al.*, 1992).

Area 2. This area extends to the north of Area 1, near the central sector of the Gulf of Cádiz. The predominant grain size is medium to fine sand, although there is also a significant proportion of bioclastic components of finer grain-size than in Area 1. The bedforms are characterized by mass-gravity flows, probably influenced by both, the Mediterranean undercurrent and down-slope flows.

Area 3. This area occupies the central sector of the Gulf of Cádiz. The bottom morphology is very complex and it is characterized by topographic ridges and canyons, oriented perpendicularly to the margin and subhorizontal platforms at several depths in the slope. The distribution of sediment types, in consequence, is controlled by the location of the topographic irregularities in respect to the Mediterranean undercurrent and the location of submarine canyons. In the areas exposed to the Mediterranean undercurrent the predominant grain-size is fine sand, while in protected areas dominates silty sand and finer materials.

Area 4. The Area 4 occupies the northern sector of the study region. It is characterised by clayey silts and silty clays, reflecting the significant decrease in energy of the Mediterranean undercurrent. The deposits develop locally contour bodies in the vicinities of the topographic irregularities

The northwestward variation in grain-size is also reflected by a host of bedforms which also record the decrease of energy of the Mediterranean undercurrent (NELSON *et al.*, 1993). Thus, sand dunes occur in the southernmost area, large mud waves are found to the West as the grain size diminishes, and large contoured bodies are observed in the northernmost regions.

REFERENCES

- BARAZA J. and NELSON H., 1992. Clasificación y dinámica de las formas de fondo en el Golfo de Cádiz: implicaciones de la corriente profunda mediterránea en los procesos sedimentarios durante el Pliocuaternario. III Congreso Geológico de España y VIII Congreso Latinoamericano de Geología. Tomo 1: 118-122.
- GUTIERREZ-MAS J. M. and VILLANUEVA GUIMERANS P., 1986. Estudio granulométrico y geoquímico de los sedimentos recientes no consolidados de la plataforma continental próxima a Cádiz. III Seminario de Química marina. Cádiz, 28, 29 Enero 1986.
- HEEZEN B. C. and JOHNSON G. L., 1969. Mediterranean undercurrent and microphysiography west of Gibraltar. *Bulletin Institut Oceanographique*, Monaco, 67: 1-95.
- MALDONADO A., 1992. Alboran sea. *Geo-Marine Letters*, Springer International, New York, v.12 (2-3) p.61-186.
- NELSON H. *et al.*, 1993. Mediterranean undercurrent sandy contourites. Gulf of Cádiz, Spain. *Sedimentary Geology*, 82: 103-131.
- PALANQUES A. *et al.*, 1986-1987. Estudio de la materia en suspensión en el Golfo de Cádiz. *Acta Geológica Hispánica* t.21-22: 491-497.