

## THE ACTIVE DEFORMATION OF ERATOSTHENES SEAMOUNT, SOUTH OF CYPRUS

J. M. WOODSIDE<sup>1</sup>, M. K. IVANOV<sup>2</sup>, R. B. KIDD<sup>3</sup>, A. F. LIMONOV<sup>4</sup>  
and A. H. F. ROBERTSON<sup>5</sup>

<sup>1</sup> Free University, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

<sup>2</sup> Moscow State University, Russia

<sup>3</sup> University of Wales, Cardiff, United-Kingdom

<sup>4</sup> Moscow State University, Russia

<sup>5</sup> University of Edinburgh, United-Kingdom

Eratosthenes Seamount is a slightly NNE/SSW-elongated block of about 100 km x 70 km which rises over 1200 m from the seafloor about 100 km south of Cyprus. Its origin is enigmatic but it is generally thought to be an uplifted or remnant block of the African continental margin. New seismic data clearly show the seamount to be underthrusting both Cyprus to the North and Nile sediments to the South, suggesting that Eratosthenes is buckling between the colliding African and Turkish plates. The plateau forming the top of Eratosthenes is separated into different levels by faults cutting it in a roughly WSW-ENE direction. Mass movements including large slides are evident on its flanks. The western flank appears to be thrusting beneath sediments to the West; but the eastern flank shows evidence of normal faulting with rotated blocks. There also seems to be a discontinuity between the seamount and the gently north-sloping Levantine basin to the East, suggesting faulting, possibly connected with the Southeast flank of the Hecataeus Ridge and the eastern part of the Cyprus arc to the North.

Northwards underthrusting beneath Cyprus, probably since Early Miocene, has caused and is causing strong compression in southern Cyprus with both uplift and southward thrusting; and the amplification of these forces since Late Miocene time was probably the result of wedging of the Eratosthenes block beneath Cyprus, which continues to the present. Because the gravity anomaly associated with Eratosthenes can be explained largely by the topography of the seamount (there is almost no associated Bouguer anomaly), it is postulated that the structure is caused in large part by up-arching since Early Miocene time with active tectonic modification during the current stage of breakup. Counterclockwise rotation of the entire block, occurring as part of the wedging process, has resulted in underthrusting of Eratosthenes beneath the sediments to the West, the cross-cutting WSW-ENE faulting, and some rifting between the eastern flank of Eratosthenes and the NE-SW fault zone defining the edge of the Levantine basin to the east of the seamount. It is therefore a very tectonically-active feature undergoing destruction in the collision process.

G