

EVOLUTION OF TRACE METAL LEVELS AND MAGNETIC PROPERTIES IN SEDIMENTS OF THE ELEFSIS GULF, GREECE

M. SCOULLOS¹, E. FOUFA² and M. DASSENAKIS¹

¹ Chemistry Dept. Athens University, Panepistimioupoli, 105 71 Athens, Greece

² General Chemical State Lab. of Greece, An. Tsocha 16, 115 21 Athens, Greece



The small (68 km²) and shallow (max depth 33 m) Gulf of Elefsis (Fig. 1) located in the northern part of the Saronikos Gulf, close to Athens and Pireaus, has received particular attention due to its scientific and ecological importance and its relation to the economy of the country. It is connected to the rest of the Saronikos Gulf with two natural narrow and shallow channels and receives considerable amounts of industrial effluents from crude oil refineries, shipyards, steel works,

cement, food, electroplating and chemical industries, mainly in its eastern part where the town and the port of Elefsis are located. In this study an attempt is made to trace any trend in the pollution of the area which is continuously studied from our laboratory since 1974 (SCOULLOS *et al.*, 1979; SCOULLOS *et al.*, 1986; FOUFA, 1993) by using a combination of chemical partitioning of trace metals and measurements of some magnetic parameters in the sediments for two sampling periods (1988 and 1992). Sea bottom sediments were collected from three sampling stations during two oceanographic cruises on May 1988 and May 1992, using a van Veen box corer. Sediment samples were wet sieved through a 61 μ m nylon net, dried at 40°C and subjected to sequential extraction. The reagents used for this procedure were (TESSIER *et al.*, 1979) : 1 : MgCl₂ 1M, pH7; 2 : CH₃COONa, 1M, pH 5; 3 : NH₂OH.HCl 0.04M, CH₃COOH 25%; 4 : H₂O₂ 30%, 0.02M HNO₃, 85%; 5 : HF-HNO₃-HClO₄ 120°C. The sediment fractions extracted respectively broadly correspond to A : Easily exchangeable, B : Carbonates, C : Fe-Mn oxides, D : Organics, E : Detrital silicates. Metal concentrations were measured with a Perkin Elmer 2380 AAS system. The magnetic parameters measured were the following (SCOULLOS & ZERI, 1993) :

- 1. Magnetic susceptibility χ represents the ease with which a material can be magnetized. It was measured using a Bartington susceptibility meter at 0.1T and 0.47kHz.

- 2. Saturation isothermal remanent magnetization (SIRM), represents the magnetic content and is measured in a fluxgate magnetometer (Minispin, Molspin Ltd) after placing the sample in a strong d.c. magnetic field (1000 mT) at 24°C temperature

- 3. Frequency depended susceptibility $\chi_{fd}\%$, defined by the ratio $[(\chi_1 - \chi_h)/\chi_1] * 100$, (χ_1 : 0.1T, 0.47kHz - χ_h : 1T, 0.47kHz). It helps in identifying very fine grains (< 0.03 μ m).

Pb-1988						%	total	
Station	A	B	C	D	E		ug/g	
1	15	10	62	6	7		274	
2	12	12	62	6	8		191	
3	11	11	62	7	9		169	
Pb-1992						%	total	
Station	A	B	C	D	E		ug/g	
1	19	14	52	10	5		264	
2	19	15	50	10	6		230	
3	20	15	49	10	6		216	

Cu-1988						%	total	
Station	A	B	C	D	E		ug/g	
1	2	4	52	34	8		188	
2	3	5	46	36	10		111	
3	2	4	48	34	12		105	
Cu-1992						%	total	
Station	A	B	C	D	E		ug/g	
1	2	3	54	32	9		178	
2	3	4	48	33	12		123	
3	2	3	51	34	10		121	

Zn-1988						%	total	
Station	A	B	C	D	E		ug/g	
1	4	5	55	27	27		632	
2	2	6	46	31	31		332	
3	3	6	40	23	23		181	
Zn-1992						%	total	
Station	A	B	C	D	E		ug/g	
1	3	5	57	26	9		606	
2	3	3	59	20	15		346	
3	3	5	44	23	25		206	

Mn-1988						%	total	
Station	A	B	C	D	E		ug/g	
1	1	21	62	9	7		1223	
2	1	22	58	9	10		734	
3	1	25	53	0	12		701	
Mn-1992						%	total	
Station	A	B	C	D	E		ug/g	
1	1	19	63	9	8		1100	
2	1	20	59	10	10		776	
3	0.1	19	59	10	12		655	

Fig.2. Fractionation of trace metals in surface sediments

The total metal content of the sediments (Fig. 2) reveals that the sediments of the eastern part of the Gulf (st. 1) are enriched in trace metals. Increased concentrations of magnetic particles of anthropogenic origin were also observed at the same station during the 1988 sampling as it becomes clear from the high SIRM and χ and the low $\chi_{fd}\%$ values. The SIRM and χ values at station 1 were found reduced at the 1992 sampling to the levels of stations 2 and 3 (Fig. 2). This may indicate a reduction of anthropogenic inputs in the area during the period 1988-92, due to the cease of some industrial activities (such as production of iron and steel). Metal concentration levels were similar in the two sampling periods but a slight decrease of total values was observed at station 1 followed by a clear increase at stations 2 and 3. This is probably caused by remobilization and/or transport of metals from particles and sediments of the eastern part of the gulf to the western part. That means that the eastern part acts now not only as a sink but also as secondary pollution source. The sequential extraction procedure for trace metals revealed that the main fraction of the examined metals was connected with Fe-Mn oxides (Fig. 1). High proportion of Cu and Zn was found in the organic fraction whereas elevated percentage of Mn and Pb was connected with carbonates. The percentage of metals held into the alumino-silicate lattice was rather limited and only for Zn exceeded 20%. Significant differentiations in metal partitioning between the two samplings were not observed. From Fig. 3 becomes clear that the variations of SIRM and χ were similar with the corresponding variations of trace metal concentrations.




Fig.3. Magnetic measurements in surface sediments




Fig.4. Variations of metal concentrations and magnetic measurements

REFERENCES

- M.J. SCOULLOS, F. OLDFIELD & R. THOMPSON, 1979. *Mar. Poll. Bull.* 10 : 287-91
 M.J. SCOULLOS & F. OLDFIELD. 1986. *Mar. Chem.* 18 : 249-68
 E. FOUFA, 1993. Heavy metals-magnetic measurements of Saronikos Gulf. MSc Thesis
 A. TESSIER, P.G. GAMBELL & M. BISSON, 1979. *Anal. Chem.* 51 : 844.
 M.I. SCOULLOS & C. ZERI, 1993. *Oceanol. Acta*, 16,1 : 53-61.

Rapp. Comm. int. Mer Médit., 34, (1995).