


GEOCHEMICAL ASPECTS OF A GULF INFLUENCED BY ANTHROPOGENIC ACTIVITIES (THERMAIKOS GULF, N.W. AEGEAN SEA)

F. VOUTSINOY-TALIADOURI, E. GEORGAKOPOULOU-GREGORIADOU & R. PSYLLIDOU-GIOURANOVITS

National Centre for Marine Research, 166 04 Hellinikon, Hellas

Thermaikos Gulf, in the N.W. Aegean Sea, receives about 120,000 m³/day of untreated sewage water from the city of Thessaloniki, with a population of 1,200,000 inhabitants. Also, an amount of about 25,000 m³/day of treated or partially treated industrial effluents is released on the northwestern coast of the bay, where the industrial zone is located. On the west side of Thermaikos Gulf, the rivers Axios, Loudias and Aliakmon



release important amounts of some metals to the sea. The main interest of the present study is to assess the environmental state of the region, as far as the heavy metal levels is concerned, after the removal of the 30% of the sewage discharges via the newly constructed outfall. It is also to compare the status of the marine environment 20 years after the first investigations.

Surface sediment samples recovered from Thermaikos Gulf during 1993 over a grid of 24 stations were examined for grain composition, organic carbon and the metals Fe, Cr, Ni, Mn, Zn, Co, Cu and Pb. The samples were taken using a 0.1m² van Veen grab. The extraction of the metals was achieved with 2N HCl and the determination of the metal content in the leachates was performed on a Perkin-Elmer 305B A. A. S. (SATS-MADJIS & VOUTSINOY-

TALIADOURI, 1981). The particle size composition was estimated according to BUCHANAN's technique (1971). The organic carbon was obtained according to GAUDETTE *et al.*, 1974. Analyses were performed in triplicate. The reliability of the whole process had been ascertained in Intercalibration Exercises (I.A.E.A., 1978). The analyses indicated the following average standard deviations and coefficient of variations: Fe (%) 0.85, 4.4; Mn (mg/kg) 38, 5.0; Zn (mg/kg) 4.3, 7.2; Cr (mg/kg) 3.8, 3.4; Ni (mg/kg) 3.3, 4.1; Co (mg/kg) 0.9, 8.2; Cu (mg/kg) 1.1, 4.8; Pb (mg/kg) 0.8, 6.7.

The cold dilute HCl extraction method is chosen because it will release both inorganic and organic associated non-residual heavy metals from sediments without materially affecting the silicate matrix (DUINKER *et al.*, 1974). The analysis of the non-residual (non-lattice held) elements will often yield more data on the extent of heavy metal pollution than will that of the total sediment which include the residual or non-polluted fraction. Non-residual heavy metals are not part of the silicate matrix and have been incorporated into the sediment from aqueous solution by processes such as adsorption and organic complexation, i.e. non-residual heavy metals include those originating from polluted waters (CHESTER & VOUTSINOY, 1981). The analyses of the data reveals that most of the Thermaikos Gulf seafloor is covered by fine sediment, which is derived mainly from the Rivers Axios, Loudias and Aliakmon and containing relatively high amounts of heavy metals. The elements studied can be divided into two categories: the ones derived mainly from anthropogenic activities and the others which are probably depended on natural geochemical processes. More specifically, the main sources of organic carbon and Cu, Pb, Zn, Cr (Fig. 1) are the Thessaloniki sewage and industrial outfalls and the Axios river. Nickel, Co, Fe and Mn have mostly a natural origin, being derived from the weathering of mafic and ultra-mafic rocks, largely extended on the adjacent land (VOUTSINOY-TALIADOURI & VARNAVAS, in press).

Table 1: Heavy metal concentrations of the present study, together with those reported in the past (10 and 20 years ago).

	Fe %	Cr ppm	Ni ppm	Mn ppm	Zn ppm	Co ppm	Cu ppm	Pb ppm
Present Study	1.94-3.14	121-294	60-224	463-1935	73-203	13-30	17-51	20-150
Study of '85*	-	66-390	55-290	215-1340	32-1610	14-37	8-170	11-330
Study of '75**	-	102-353	52-240	347-2050	45-280	12-42	11-82	13-230

* VOUTSINOY-TALIADOURI, F. & LEONARIS, S.N.

** CHESTER, R. & VOUTSINOY, F.G. After the modification with a factor which equilibrates the two extraction techniques.

Table 1 gives the results obtained from this study, as well as results obtained with the same methodology 10 and 20 years ago. As it can be seen, the heavy metal ranges do not have changed during the 20 last years. The relatively higher maximum values of the non-residual heavy metals (Cr, Zn, Cu and Pb) observed in the study of '85 are mainly due to the fact that the sampling stations in that study were closer to the pollution sources.

REFERENCES

- BUCHANAN J.B., 1971. Sediments, in: Methods for the study of marine benthos, ed. by N.A. Holme & A. McIntyre, IBP Handbook n°16, Oxford University Press, Oxford, pp. 35-39.
- CHESTER R. & VOUTSINOY F.G., 1981. The Initial Assessment of Trace metal Pollution in Coastal Sediments. *Mar. Pollut. Bull.*, 12: 84-91.
- DUINKER J.C., VAN ECK G.T.M. & NOLTIN, R.F., 1974. On the behaviour of copper, zinc, iron and manganese and evidence for mobilization processes in the Dutch Wadden Sea. *Neth. J. Sea Res.*, 8: 214-239.
- GAUDETTE H., FLIGHT W., TONES I. & FOLGER D., 1974. An inexpensive titration method for the determination of organic carbon in recent sediments. *J. Sedim. Petrol.*, 44: 249-253.
- IAEA., 1978. Intercalibration of analytical methods on marine environmental samples. Progress Report n°18.
- SAISMADJIS J. & VOUTSINOY-TALIADOURI F., 1981. Determination of trace metals at concentrations above the linear calibration range by electrothermal atomic absorption spectrometry. *Anal. Chim. Acta*, 131: 83-90.
- VOUTSINOY-TALIADOURI F. & LEONARIS S.N., 1986. An assessment of metal Pollution in Thermaikos Gulf, Greece. *Rapp. Comm. int. mer Médit.*, 30: 2-43.
- VOUTSINOY-TALIADOURI F. & VARNAVAS S.P., 1994. Geochemical and sedimentological patterns in the Thermaikos Gulf, N.W. Aegean Sea, *Estuar., Coast. and Shelf Sc.*, in press.

Rapp. Comm. int. Mer Médit., 34, (1995).

