

GROWTH DYNAMICS OF SPRAT *SPRATTUS SPRATTUS* L. OFF BULGARIAN BLACK SEA COAST

Gueorgui DASKALOV

Institute of Fisheries, Varna, Bulgaria

Sprat shows remarkable variability in size and growth rate between years and this feature creates its specific adaptive response to changes in environment. In this study we analyse the growth of cohorts 1977 to 1990 in relation to some environmental and population characteristics. Growth was modelled on the base of monthly length-at-age data by fitting the von Bertalanffy growth function (VBGF). Growth performance index : $\phi' = \log_{10}k + 2\log_{10}L_{\infty}$ (PAULY and MUNRO, 1984) together with direct length-at-age observations were used for growth comparisons. Correlation analysis was performed on growth parameters and environmental indices (Table 1.)

Table 1. Correlation matrix of growth and environmental parameters: ϕ' -growth performance; L1,L2-length-at-age 1 and 2 years; L_{∞} -mean length; $\Delta L1, \Delta L2$ -annual increment at age 1 and 2; c.f.-condition factor; R-recruitment; N1+,B1+-stock numbers and biomass at age 1 and older; F-fishing mortality; S.I.-spawning intensity¹; Zoo, Ph-zoo- and phytoplankton biomass; PO₄, C.W.- phosphate concentration, water temperature and winter cold² in the N-W Black Sea. Significance levels: * - p = 0.05; # - p = 0.01

year	ϕ'	L1	L2	L_{∞}	$\Delta L1$	$\Delta L2$	c.f.	R	N1+	B1+	F	S.I.	Zoo	Ph	PO ₄	T ^o
ϕ'	0.09															
L1	-0.91#	-0.27														
L2	-0.87#	0.16	0.77#													
L_{∞}	-0.89#	0.15	0.78#	1.00#												
$\Delta L1$	0.52	0.79#	-0.73#	-0.28	-0.31											
$\Delta L2$	0.71#	0.04	-0.82#	-0.69#	-0.72#	0.55*										
c.f.	-0.22	0.53	0.27	0.34	0.34	0.30	-0.51									
R	-0.67#	-0.09	0.68#	0.43	0.45	-0.56*	-0.65*	0.03								
N1+	0.83#	-0.48	0.84#	0.66*	0.68#	-0.76#	-0.56*	-0.37	0.52							
B1+	-0.91#	-0.33	0.88#	0.70#	0.73#	-0.70#	-0.67#	-0.19	0.62*	0.97#						
F	-0.02	0.52	-0.15	0.22	0.18	0.55*	0.25	0.66*	-0.10	-0.36	-0.32					
S.I.	0.43	0.70#	-0.44	0.03	-0.02	0.70#	0.23	0.65	-0.43	-0.73*	-0.67*	0.69*				
Zoo	-0.75#	-0.08	0.68#	0.83#	0.82#	-0.40	-0.52	0.19	0.30	0.69#	0.68#	0.13	-0.02			
Ph	-0.58*	-0.23	0.41	0.56*	0.58*	-0.36	-0.38	-0.20	0.26	0.58*	0.58*	-0.32	-0.46	0.50		
PO ₄	-0.46	-0.39	0.44	0.61*	0.57	-0.35	0.31	-0.22	-0.06	0.38	0.34	0.00	-0.22	0.73#	0.71#	
T ^o	0.70	0.78*	-0.81*	-0.56	-0.66	0.88#	0.71*	0.14	-0.25	-0.80*	-0.74*	0.70	0.81*	-0.65	-0.50	-0.51
C.W.	-0.69	-0.77	0.48	0.10	0.13	-0.63	-0.06	-0.45	0.37	0.62	0.54	-0.42	-0.90#	0.24	0.52	0.42

1 As a relative index of interannual variability of the spawning intensity was used the average percentage of fishes with ovaries in maturity stages IV and V during the peak spawning season: November - January.

2 Winter conditions are important because of the positive effect of the winter convection (which is particularly intensive in cold and windy winters) on bioproductivity.

An intensification of sprat fishery started in the mid 70's on the base of rising stock abundance, due to outstanding "eutrophic" productivity of the Black Sea and reduced predatory press. After 1980, sprat biomass being hard exploited, dropped down in Bulgarian waters (PRODANOV and DASKALOV, 1992). In terms of growth, the period 1977-1993 is characterized by decrease in size and relative increase in growth rate till

year	L _{oo}	k	ϕ'	$\Delta L1$	L	c.f.
1977	12.62	0.329	1.719	0.89	10.59	-
1978	30.73	0.042	1.598	0.89	10.13	0.58
1979	14.30	0.271	1.744	1.31	10.17	0.553
1980	16.85	0.145	1.615	1.04	10.67	0.587
1981	12.41	0.594	1.961	1.79	10.40	0.614
1982	12.80	0.427	1.845	1.48	10.23	0.616
1983	13.21	0.344	1.778	1.37	10.10	0.585
1984	12.02	0.544	1.895	1.44	10.27	0.588
1985	13.50	0.282	1.711	1.27	9.80	0.596
1986	12.65	0.404	1.811	1.49	9.80	0.576
1987	26.03	0.069	1.670	1.19	9.27	0.581
1988	19.36	0.129	1.684	1.39	9.57	0.554
1989	15.34	0.230	1.733	1.57	9.06	0.568
1990	12.27	0.399	1.770	1.45	9.10	0.593

Table 2. Growth parameters of sprat




Fig. Growth curves for average cohorts 1977-80, 1981-83, 1984-86, 1987-90.

The changes in growth of sprat can be associated mainly with the gradual reducing of the standing stock under intensive exploitation. After 1986, planctivorous invertebrates (especially the ctenophore *Mnemiopsis* sp.) become dominant in the pelagic community. Competition on food with fish larvae could be one possible explanation of the decrease in growth in the last years.

ACKNOWLEDGEMENTS. the present study was partly supported by the Research Support Scheme of the Central European University, no:1821041191-92.

REFERENCES

- PAULY, D. and J.L. MUNRO, 1984. *Fishbyte* 2(1): 21
 PRODANOV, K. and G. DASKALOV, 1992. *Rapp. Comm. int. Mer Médit.* 33: 305
Rapp. Comm. int. Mer Médit., 34, (1995).