

ADRIATIC SEICHE DECAY AND ENERGY LOSS THROUGH OTRANTO STRAIT

Ivana CEROVECKI¹, Mirko ORLIC² and Myrl HENDERSHOTT³

¹ Inst. Ruder Boskovic, Center for Marine Research, Bijenika 54, 41000 Zagreb, Croatia

² Geophysical Inst., Fac. of Science, Univ. of Zagreb, 41000 Zagreb, Croatia

³ Scripps Institution of Oceanography 0209, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0209, USA

A salient feature of sealevel records from the Adriatic Sea is the frequent occurrence of energetic seiches with a period of about 21 hours. Once excited by a sudden wind event, such seiches may persist for days. They may lose energy either to friction within the Adriatic, or by radiation through the Straits of Otranto into the Mediterranean. The decay caused by both mechanisms has been included in a one dimensional variable cross section shallow water model of the Adriatic. Friction is parameterized by the coefficient k appearing in the linearized frictional acceleration term $-ku/h$ (where h is the depth and u is the shallow water along-basin velocity), and radiation through Otranto is described by the coefficient a appearing in the strait boundary condition $z = auh/(\text{sqrt } gh)$ in which additionally z is sealevel and g the acceleration of gravity. Repeated runs of the model delineate the dependence of model seiche decay time on k and a shown in the accompanying figure.

Actual decay times of the fundamental longitudinal mode seiche have been determined by constructing the decay envelopes of sealevel (detided and subjected to a narrow band filter of width .0375 oph centered around the seiche frequency) from three locations (Bakar, Split and Dubrovnik) along the Croatian coast during twelve seiche episodes between 1963 and 1986, and taking into account only those portions of the envelopes which decrease exponentially in time and for which the modeled effects of along-basin winds were smaller than the error of estimation of decay time from the envelopes and the effects of across-basin winds were small. The average decay time thus obtained was 3.2 ± 0.5 days.

For the model to reproduce this observed decay time with no energy loss through Otranto, a friction coefficient of $k = .00067$ m/s was needed; to reproduce the observed decay time with no friction, a coefficient of energy transmission of $a = .064$ was required. To proceed further, an estimate of k that is independent of observations of seiche decay time is required. ORLIC *et al.* (1986) estimate $k = .00113 - .00124$ m/s by balancing surface and bottom stresses in wind driven currents in the Northern Adriatic; ORLIC (1987) estimates $k = .0006 - .00153$ m/s from the observed decay time of near-inertial oscillations in the Northern Adriatic. The corresponding values of a needed for the model to reproduce the observed seiche decay time are zero to 0.0085; these imply the loss of zero to 3.3% of the seiche energy to the Mediterranean every seiche period. Since the decay time might be as short as 2.7 days, a could be as great as .02; in this case 7.8 % of the seiche energy would be lost to the Mediterranean every seiche period.

REFERENCES

- ORLIC M., 1987. Oscillations of the inertial period on the Adriatic Sea Shelf. *Cont. Shelf. Res.*, 7: 577-598.
ORLIC M., Kuzmic M., and Vucak Z., 1986. Wind-curl currents in the Northern Adriatic and formulation of bottom friction. *Oceanol. Acta*, 9: 425-431.