

STOCK ASSESSMENT OF SPRAT *SPRATTUS SPRATTUS* L. OFF BULGARIAN BLACK SEA COAST, USING LENGTH COHORT ANALYSIS

Gueorgui DASKALOV¹ and Kamen PRODANOV²

¹ Institute of Fisheries, Varna, Bulgaria

² Institute of Oceanology, Varna, Bulgaria

First attempt is made to evaluate the dynamics of abundance and mortality of sprat stock by means of the Length Cohort Analysis (LCA) (JONES, 1981).

Length composition data for Bulgarian sprat catches was averaged generally by 3 years over the period 1945-92 and the resulting pseudo-cohorts were assumed to be under steady state conditions.

LCA was performed using ANALEN software package (CHEVAILLIER et LAUREC, 1990), where an iterative procedure for running the analysis was employed. It was also possible to estimate the fishing mortality rates per fishing gear for 1970-92, when separated statistics for coastal (trapnets, beach seines) and trawl fisheries were available. Sensitivity analysis was taken into account, for the optimal choice of parameters.

It was not possible to estimate directly the von Bertalanffy's parameters, because of lack of satisfactory data covering all the period studied. Even though such estimates were available for 1977-93 (DASKALOV, this vol.), we did not obtain reasonable results using these values. The procedure suggested by JONES (1990) gives closest results with previous studies (DASKALOV, 1993).

Table. A : Parameters used in analysis. B : Basic results.

R= recruitment in 106 numbers, \bar{B} = mean biomass in tons, F = average fishing mortality rate

years	A			B		
	k(Loo=16)	M	F+	R	\bar{B}	F
45-49	0.156	1.2	0.014	4449.8	7575.1	0.108
56-59	0.155	1.2	0.026	9877.7	17634.0	0.089
60-62	0.164	1.2	0.013	55244.1	133686.2	0.007
63-65	0.188	1.2	0.011	12224.2	26035.4	0.029
66-68	0.239	1.2	0.006	12998.0	32174.3	0.014
69-71	0.239	1.2	0.014	9354.6	21751.7	0.046
72-74	0.265	1.2	0.024	311150.0	79686.2	0.024
75-77	0.241	0.95	0.041	29197.2	113437.9	0.028
78-80	0.185	0.95	0.088	46438.5	176661.6	0.039
81-83	0.314	0.95	0.06	16373.7	58990.0	0.146
84-86	0.267	0.95	0.04	13682.3	39556.5	0.237
87-89	0.237	0.95	0.028	13553.4	37006.7	0.135
90-92	0.217	0.95	0.009	7766.65	23495.6	0.079

The average value Loo = 16cm (1977-92) was assumed for all the period and then k was found :

$$k = \ln((Loo-11)/(Loo-12))$$

where 11 and 12 were respective lengths-at-age 1 and 2 years (Table, A). Terminal length group (+) was chosen to be 12 or 11.5 cm (> 75%Loo). Natural mortality rate was assumed - 0.95 for 1975-92 and 1.2 for 1945-74 - the last one for the purpose of reflecting the higher predation in that period (STOYANOV, 1966, IVANOV and BEVERTON, 1985). Terminal fishing mortality rates were obtained according to DASKALOV's VPA estimates (1993). A functional regression built between catches (C) and fishing mortality rates (F) with coastal gears for 1975-92 was used to determine F+ values before 1975.




Fig. Sprat mean biomass (B : columns) and catches (C : line) in tons : 1945-92

The evolution of the sprat stock state could be devised into three main stages (Table, B, Fig.). In the years up to mid 70's, stock biomass remains relatively low in relation with a strong predatory press. As an exception, we observe the period 1960-62 where higher abundance is probably resulting from the combination of stable recruitment and favourable environment at the end of 50's. The second stage is characterised by a very strong increase of stock biomass and fishing from mid 70's to mid 80's. Such an "explosion" could be related with the extinction of top predators in late 60's and early 70's, and the rise of the sea trophic level due to progressive eutrophication. The combined action of two factors explains the decrease in sprat biomass after the late 70's. In the beginning, the high nutrient abundance resulted in amelioration of the trophic base, but soon the outstanding eutrophication created different negative effects, like hypoxia and increasing domination of gelatinous megaloplankton, which is feeding on fish eggs and larvae and competes planktivorous fish on food. The second factor is obviously the fishing effort remaining too high at the same time when the standing stock is decreasing.

REFERENCES

- CHEVAILLIER, P. et LAUREC, A. 1990. *FAO Doc. Tech.Pêches*, 101(4) : 124 pp.
 DASKALOV, G. 1993. *Rapport de DEA, Univ. Aix-Marseille II, OSU(COM)*
 IVANOV, L and BEVERTON, R.S.H., 1985. *FAO Stud.Rev.*, 60 : 135 pp.
 JONES, R., 1981, *FAO Fish.Circ.*, 734 : 57 pp.
 JONES, R., 1990, *J. Cons. int. Explor. Mer*, 46 : 130-139
 STOYANOV, S., 1966, *Izv.Nauchn.Inst.Rib.Stop.Okeanogr.*, Varna, 6 : 21-48

