

OCCURRENCE OF CORIS JULIS(LINNAEUS, 1758) LARVAE IN THE SOUTHERN TUSCAN ARCHIPELAGO

Sartini M. ¹, Belcari P. ^{2*}

¹ Aplysia p.s.c.r.l., Livorno, Italia - aplysia1@libero.it

² Dipartimento di Scienze dell'Uomo e dell'Ambiente, Università di Pisa, Italia

Abstract

Distribution, abundance and size structure of *Coris julis* larvae were analysed. Data were collected during three ichthyoplankton cruises carried out in the southern Tuscan Archipelago (northern Tyrrhenian Sea) in 1999 and 2000. *C. julis* larvae were only collected during June, showing a wide distribution on the continental shelf.

Keywords: Ichthyoplankton, larvae, distribution, abundance, northern Tyrrhenian Sea

Introduction

The rainbow wrasse, *Coris julis* (Labridae), is the most common and abundant wrasse in the Mediterranean, especially in its western basin. The species lives on Posidonia beds as well as rocky and mixed bottoms, from few meters to 120-150 m depth, but it is more frequent in waters shallower than 50 m (1). It is a by-catch of the inshore trawl and of trap and angling fisheries. Some biological aspects of *C. julis* have not yet been deeply investigated, probably because of the lack of commercial value of the species. The aim of this study was to give some information on the abundance, distribution and size structure of *C. julis* larvae collected in the southern Tuscan Archipelago.

Material and methods

In the southern Tuscan Archipelago, northern Tyrrhenian Sea, three ichthyoplankton cruises were carried out in March 1999 and March and June 2000 (2). During each cruise, a minimum of 60 stations were occupied, collecting a total of 210 samples. Larvae were sampled by means of oblique hauls, from near the bottom to the surface. A Bongo net with 60 cm mouth diameter and 500 μ m mesh size was used. Sampling methodology followed standard procedures (3). Larvae of *C. julis* were identified according to morphological, morphometric and pigmentation features (4). In addition, the standard length (SL, to the nearest 0.1 mm) was measured for each specimen. Larval abundance was standardised to the volume of filtered water at each station and was expressed as number of larvae/10 m² of sea surface (3).

Results and discussion

In the ichthyoplankton survey carried out in June 2000, 113 larvae of *C. julis* were collected, whereas during the cruises of March 1999 and March 2000 no *C. julis* larvae were collected. This is in agreement with existing knowledge of the reproductive period of the species which occurs in late spring-early summer (5).

C. julis larvae presented a wide spatial distribution (Fig. 1). The majority of stations at which they were found were localized between 50 and 200 m, in accordance with the ecological characteristics of the adults. The presence of larvae at stations deeper than 200 m could be related to advection by sea currents in the study area (6).

The maximum abundance of *C. julis* (72 larvae/10 m²) was recorded below the Island of Elba, near the Island of Pianosa; the minimum abundance (4 larvae/10 m²) was found at stations located near the Islands of Giglio and Giannutri (Fig. 1).




Fig. 1. Distribution and abundance of *C. julis* larvae, expressed as number/10 m² of sea surface, estimated for the ichthyoplankton survey of June 2000.

The size of the larvae ranged from 2.04 to 6.56 mm SL (Fig. 2). Most specimens (about 76%) were between 2.5 and 4 mm SL, with a mode at 3.5 mm SL.




Fig. 2. Size frequency distribution of *C. julis* larvae collected during the ichthyoplankton survey of June 2000.

References

- 1-Quignard J.P., Pras A., 1986. Labridae. Pp. 919-942. In: Whitehead P.J.P., Bauchot M.-L., Hureau J.C., Nielsen J., Tortonese E. (eds.), Fishes of the North-eastern Atlantic and the Mediterranean. UNESCO, Paris, vol. 2.
- 2-Sartini M., 2003. Ittioplankton dell'Arcipelago Toscano meridionale. Ph. D. Thesis, University of Pisa, 233 p.
- 3-Smith P.E., Richardson S., 1977. Standard techniques for pelagic fish eggs and larval surveys. FAO Fish. Tech. Pap., No. 175, 100 p.
- 4-Spartà A., 1933. Contributo alla conoscenza di uova e larve nei Labridi. IV. Uova e larve di *Julis G. G. Risso* e *Julis pavo* ottenute da fecondazione artificiale, e stadi post-embryonali raccolti nel plancton. R. Com. Talassogr. Ital., Mem. 203, 18 p.
- 5-Fiorentino F., 1999. *Coris julis*. In: Relini, G., Bertrand, J., Zamboni, A. (eds.), Synthesis of the Knowledge on Bottom Fishery Resources in Central Mediterranean (Italy and Corsica). Biol. Mar. Medit., 6 (suppl. 1): 222-228.
- 6-Elliott A.J., 1979. Low-frequency sea level and current fluctuation along the coast of Northwest Italy. J. Geophys. Res., 84: 3752-3760.