

²¹⁰PO AND ²¹⁰PB CONCENTRATIONS, FLUXES, PARTICLE SETTLING VELOCITIES, AND ORGANIC CARBON AT THE DYFAMED SITE, NORTHWESTERN MEDITERRANEAN

Gillian Stewart ^{1*}, J. Kirk Cochran ¹, Pere Masqué ², Robert Armstrong ¹, Juan-Carlos Miquel ^{3, 4}, Alessia Rodriguez, Scott Fowler ³, Michael Peterson ⁵, Nicholas Fisher ¹

¹ Marine Sciences Research Center, Stony Brook University, USA - * gstewart@ic.sunysb.edu, kcochran@notes.cc.sunysb.edu, nfisher@notes.cc.sunysb.edu rarmstrong@notes.cc.sunysb.edu

² Institut de Ciència i Tecnologia Ambientals, Universitat Autònoma de Barcelona, Bellaterra, Spain - Pere.Masque@uab.es

³ IAEA Marine Environment Laboratory, Monaco - J.C.Miquel@iaea.org, S.Fowler@iaea.org

⁴ Dipartimento per lo Studio del Territorio e delle sue Risorse, University of Genoa, Italy and IAEA, Monaco - A.M.Rodriguez@iaea.org

⁵ University of Washington, USA - mlplmp@u.washington.edu

Abstract

Novel settling-velocity sediment traps and an elutriator were employed along with traditional techniques to explore the characteristics of sinking and suspended material at the DYFAMED site. This is the first time that ²¹⁰Po, ²¹⁰Pb, and other characteristics have been examined in particulate samples separated by sinking velocity and may further elucidate the composition of particulate flux. Preliminary results suggest that ²¹⁰Po activities differ among particles sinking at different rates. These measurements will allow us to compare the efficiency of ²¹⁰Po and ²³⁴Th as flux tracers, with an emphasis on the potential usefulness of ²¹⁰Po as a POC flux tracer.

Keywords: Polonium-210, Lead-210, Flux, Organic Carbon, Thorium

Traditionally, sinking and suspended particles in the ocean have been separated by size fractions using filters and meshes (e.g. 1). Size alone, however, does not determine the rate at which particles sink. Equally important are the composition of the material and its density. In an attempt to establish some relationship between composition and sinking velocity, we collected particles at the DYFAMED site and separated them based solely on their settling rates. The samples were divided using an elutriator and a specially designed sediment trap into six sinking velocity ranges between <10 and >230m per day; each fraction was analyzed for total and organic carbon, ballast minerals, protein, lipids, pigments, and the radionuclides ²¹⁰Po, ²¹⁰Pb, and ²³⁴Th.

Previously, the ratio of polonium to thorium has been used to decipher the organic and biogenic silica content of particles in the Southern Ocean (2). In the context of MEDFLUX, we are interested in the relationship between organic content, ballast content, and sinking flux. Potentially the amount of ballast, or dense mineral content, in a particle may determine not only the sinking rate, but also the extent to which organic matter is preserved as it is transported to depth (3,4). The preservation of organic carbon by ballast below the mixed layer may successfully sequester carbon for thousands of years, and thus warrants further investigation for the accurate modeling of the global carbon cycle and the biological pump.

Specifically, we are looking for a correlation between the ²¹⁰Po/²¹⁰Pb or the ²¹⁰Po/²³⁴Th ratio and the organic content of sinking matter. We predict that the ratio of polonium to lead and/or thorium will reflect the ratio of organic matter to ballast as ²¹⁰Po is known to associate with labile cellular material (5) and ²¹⁰Pb and ²³⁴Th are linked with more refractory organic material or inorganic substances such as sediment, atmospheric dust, and biogenic minerals (6,7).

Preliminary data suggest that the polonium activity differs in particles with differing sinking velocities (Fig 1). Over 52% of the to-

tal polonium flux was found within the fastest sinking particles but, when normalized to mass, the particles sinking between 29 and 58 m/day had the highest ²¹⁰Po activity. This result indicates that these particles contain either the highest organic content, highest protein content, or concentrate polonium for some other reason. The nature of these particles will be more fully known when the other analyses (total and organic carbon, Al, Ti, Si, etc.) are complete. Results of these investigations as well as the lead and thorium activities will be presented.

References

- 1 - Harvey H.R. and J.R. Johnston, 1995. Lipid-composition and flux of sinking and size-fractionated particles in Chesapeake-Bay. *Organic Geochemistry*, 23: 751-764.
- 2 - Friedrich J. and M.M. Rutgers van der Loeff, 2002. A two tracer (²¹⁰Po-²³⁴Th) approach to distinguish organic carbon and biogenic silica export flux in the Antarctic Circumpolar Current. *Deep Sea Research*, 49: 339-354.
- 3 - Mayer, L.M., 1994. Surface area control of organic carbon accumulation in continental shelf sediment. *Geochimica Cosmochimica Acta*, 58: 1271-1284.
- 4 - Armstrong, R.A., C. Lee, J.I. Hedges, S. Honjo, and S. Wakeham, 2002. A new, mechanistic model for organic carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals. *Deep Sea Research*, 49: 219-236.
- 5 - Stewart G.M. and N.S. Fisher, 2003a. Experimental Studies on the accumulation of polonium-210 by marine phytoplankton. *Limnology and Oceanography*, 48(3): 1193-1201.
- 6 - Nozaki, Y., F. Dobashi, Y.Kato, Y. Yamamoto, 1998. Distribution of Ra isotopes and the ²¹⁰Pb and ²¹⁰Po balance in surface waters of the mid Northern Hemisphere. *Deep Sea Research*, 45: 1263-1284.
- 7 - Bacon M.P. and R.F. Anderson, 1982. Distribution of thorium isotopes between dissolved and particulate forms in the deep sea. *Journal of Geophysical Research*, 46: 1293-1299.

Activity (Bq/g) per Elutriator Fraction




Fig. 1. The preliminary ²¹⁰Po activity (Bq/g) in each of the elutriator fractions. The numbers below each column represent the minimum settling velocity in meters per day of particles in that fraction. The first column includes particles sinking faster than 230 m/day while the final column includes all particles that sink slower than 29 m/day.