

IN-SITU TEMPERATURE MEASUREMENTS AND GEOCHEMICAL POREWATER DATA SUGGEST HIGHLY DYNAMIC RECENT ACTIVITY OF ISIS MUD VOLCANO, EASTERN MEDITERRANEAN SEA

Tomas Feseker¹ *, Jean-Paul Foucher¹, Anke Dählmann³ and François Harmegnies¹

¹ Ifremer Centre de Brest, Plouzané, France - tomas.feseker@ifremer.fr

² Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

³ University of Utrecht, The Netherlands

Abstract

In-situ temperature measurements and geochemical porewater data from Isis mud volcano were obtained during the Nautinil and Mimes cruise in 2003 and 2004, respectively. The observations show a high level of recent activity and reveal significant changes in both the temperature and chlorinity profiles from the center of the mud volcano. Coupled geothermal and geochemical models demonstrate that the changes cannot be explained by simple variability of upward fluid flow and suggest the presence of an additional, effective cooling and mixing process, which could be related to gas ebullition or seawater circulation.

Keywords : *Mud Volcanoes, Eastern Mediterranean, Models, Temperature, Geochemistry.*

The Nile deep-sea fan is the most important sedimentary accumulation in the eastern Mediterranean. Two geophysical mapping campaigns conducted in 1998 (Prismed II) and 2000 (FaniI) resulted in a first detailed morphostructural description of this deep-sea fan [1-4] and led to the discovery of several circular and sub-circular sedimentary structures on the upper slope of the eastern province. Up to a few kilometers in diameter and generally showing a low relief of a few tens of meters, these mud volcanoes or so-called 'mud pies' have been described as surface expressions of deep-seated gas chimneys [4]. Isis mud volcano (Isis MV) is located at a water depth of approximately 1020 m. In contrast to the more complex structure of the neighboring Osiris and Amon mud volcanoes, Isis MV shows a very distinct camembert-shaped morphology with relatively steep flanks and a gently domed top elevated between 20 and 40 m above the surrounding seafloor.

Detailed geothermal and geochemical investigations of Isis MV have been carried out during the Nautinil (2003) and Mimes (2004) cruises within the framework of the Euromargins/Mediflux project [5]. Sediment temperatures of more than 40 °C at 10 m below the seafloor at the center of the mud volcano indicate an exceptionally high level of activity. Rapidly decreasing temperature gradients away from the center support the hypothesis of an axisymmetric dynamic functioning of Isis MV. At the center, pore water profiles show a rapid decrease of chlorinity within the uppermost meter of the sediments, whereas the chlorinity of cores taken at short distances away is equal to bottom water values.

The data obtained during the two cruises reveal significant changes in both the sediment temperature profiles and the geochemical porewater profiles of cores from the center of the mud volcano. In 2003, *in-situ* measurements revealed a slightly convex-upward temperature profile with a gradient of up to 2.78 °C/m at depths between 2 and 7 m below the seafloor. When the measurements were repeated at the same location in 2004, the sediment temperatures had decreased by more than 5 °C at approximately 7 m below the seafloor and the previously observed convex-upward curvature of the profiles had disappeared. The temperature gradient did not exceed 2.3 °C/m, corresponding to a decrease in heat flow of about 17 percent with respect to the measurements in 2003. The chlorinity profile of the piston core that was sampled during the Nautinil cruise in 2003 showed a rapid decrease from the seawater concentration of about 608 mmol/kg to less than 170 mmol/kg less than 1 m below the seafloor. An equally sharp transition was observed in the chlorinity profiles of the cores that were sampled during the Mimes cruise in 2004. However, at this time the chlorinity decrease was shifted to a sediment depth of 0.5 to 1 m and the transition zone was overlain by normal seawater concentration in the upper part of the cores.

A one-dimensional coupled model of heat transfer and solute transport in the upper sediment column has been developed to analyze the observed changes and to characterize the associated processes. The model results indicate that the variations in the temperature and chlorinity profiles exceed the effects that may be expected from simple heat conduction and solute diffusion. Neither the rapid temperature decrease nor the downward shift of the chlorinity step observed between 2003 and 2004 can be explained by varying rates of upward fluid flow. Instead, the model suggests the presence of an additional process that leads to cooling of the sediment

column and flushing of the upper meter of the sediment with seawater at the same time. In order to identify this process, the role of gas ebullition and the possibility of seawater circulation within the upper meters of the sediment column are currently investigated. New insights into the dynamics of the activity of Isis MV are expected from the Bionil cruise in 2006.

References

- 1 - Mascle, J., Zitter, T., Bellaiche, G., Droz, L., Gaullier, V., Loncke, L., and Prismed Scientific Party (2001): The Nile deep sea fan: preliminary results from a swath bathymetry survey. *Marine and Petroleum Geology*, 18, 471-477.
- 2 - Sardou, O. and Mascle, J. (2003): Cartography by multibeam echosounder of the Nile deep-sea Fan and surrounding areas. Special publication CIESM, Monaco.
- 3 - Loncke, L., Gaullier, V., Mascle, J., and Vendeville, B. (2002): Shallow structure of the Nile deep-sea fan: interactions between structural heritage and salt tectonics: consequences on sedimentary dispersal. *In: CIESM (2002): Turbidite systems and deep-sea fans of the Mediterranean and the Black seas. CIESM Workshop Series, 17, Monaco.*
- 4 - Loncke, L., Mascle, J., and FaniI Scientific Parties (2004): Mud-volcanoes, gas chimneys, pockmarks and mounds in the Nile deep-sea fan (Eastern Mediterranean): geophysical evidences. *Marine and Petroleum Geology*, 21, 669-689.
- 5 - Feseker, T., Dählmann, A., and Foucher, J.-P. (2006): Thermal and geochemical evidence for episodic mud eruptions at a mud volcano? The Isis mud volcano case. *In: CIESM (2006): Fluid seepage / mud volcanism in the Mediterranean and adjacent domains. CIESM Workshop Series, 29, Monaco.*