

MESURHO : A HIGH FREQUENCY OCEANOGRAPHIC BUOY AT THE RHONE RIVER MOUTH

J. F. Cadiou^{1*}, M. Repecaud¹, M. Arnaud², C. Rabouille³, P. Raimbaud⁴, O. Radakovitch⁵, S. Meule⁵ and P. Gaufrès⁶

¹ Institut Français de Recherche pour l'Exploitation de la Mer (IFREMER), BP 330, 83507 La Seyne sur Mer – France - jfcadiou@ifremer.fr

² Institut de Radioprotection et de Sureté Nucléaire (IRSN), BP 330, 83507 La Seyne sur Mer – France

³ Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Avenue de la Terrasse 91198 Gif-sur-Yvette – France

⁴ Centre d'Océanologie de Marseille (COM), 163 avenue de Luminy 13288 MARSEILLE – France

⁵ CEREGE, Europôle de l'Arbois 13545 Aix-en-Provence – France

⁶ Centre d'Etudes Techniques Maritimes Et Fluviales (CETMEF) 2, Bd Président Kennedy - 13097 Aix-en-Provence – France

Abstract

As the current main source of continental fresh water discharge in the Mediterranean Sea, the Rhone river inputs are a key forcing for ecosystems of the Gulf of Lions in the northern part of the western basin. An oceanographic buoy has been installed at the Rhone river mouth in June 2009 for high frequency multi-parameter measurements with bottom, water column and air sensors. Data are sent in near real time towards an onshore data center.

Keywords: Rhone Delta, Coastal Systems, Gulf Of Lions, Instruments And Techniques, River Input

With an average flow of $1700 \text{ m}^3 \cdot \text{s}^{-1}$, the Rhone river is currently the main source of fresh continental water inputs in the Mediterranean Sea. Draining an industrialized catchment, the Rhone river is therefore a major forcing for the western Mediterranean Sea and the Gulf of Lions ecosystems. Its contribution to riverine inputs in the Gulf of Lions is estimated to be more than 85% for water and 80% for suspended particulate matter (SPM) [1, 2]. Understanding and modeling the functioning of the coastal system in this area requires a good knowledge of the quantity of water, SPM, nutrients and pollutants brought to the sea by the Rhone river.

In order to complete the multi-parameter measurements acquired at the SORA^[a] monitoring station located in Arles 50 km upstream from the sea a new instrumented platform was installed at the Rhone river mouth. It will help studying the complex processes occurring in the fresh/saline water transition zone. In the Mediterranean area and especially in the Gulf of Lions, extreme events such as floods and storms are known to play a key role in the ecosystem dynamic. To observe these events, a high frequency observation system has been designed. Coupling high frequency measurement with near real time data transmission will allow detecting an event and triggering additional sampling. It will also provide the mean to periodically check the instrument functioning.

In June 2009, the East Roustan buoy ($43^\circ 19.2 \text{ N}$, $4^\circ 52 \text{ E}$, 20 meters water depth) which is one of the two marker buoys of the river prodelta has been replaced by a new platform equipped with oceanographic instrumentation. This new system consists of a buoyancy beacon (fig. 1) which is suitable for small range tide environment [3]. Compared to a chain mooring buoy, this beacon allows simple cable connections between the surface platform and the sea bottom instruments. Bottom sensors get energy from a solar panel unit and are connected to an electronic management and communication device installed in the top part of the buoy.

Initially, the buoy was equipped with two multi-parameter sensors (one below the subsurface and the other close to the sea bottom), a weather station, a photosynthetic active radiation sensor (PAR) and an acoustic current Doppler profiler (ADCP). The acquisition and transmission of periodic data to the Coriolis *in situ* data center (www.coriolis.eu.org) via the GSM network is operational and an interface for online consultation of the data is available. In a second phase, further instruments will be added, including a nutrient sensor, a high resolution altimeter and a benthic station equipped with oxygen microelectrodes for sediment remineralization studies.

This new observation system, Mesurho, is a part of the coastal operational oceanography *in situ* network to be developed in the north western Mediterranean. The collected data will be used among others by the Previmer forecasting and analysis system (www.previmer.org).

Mesurho project has received support from the Ifremer MEDICIS program and the French national research agency (ANR) through the EXTREMA and CHACCRA projects. It has received funding support from the Provence Alpes Côte d'Azur Region.

^[a] SORA : Station d'Observation du Rhône à Arles




Fig. 1. Mesurho buoy instruments

References

- 1 - Evaluation des flux polluants à la Méditerranée (hors Rhône) – SIEE 2003 – Agence de l'Eau Rhône Méditerranée & Corse
- 2 - Durrieu de Madron X., Abassi A., Heussner S., Monaco A, Aloisi J.C., Radakovitch O., Giresse P. Buscaill R., Kerhervé P., 2000. "Particulate matter and organic carbon budgets for the Gulf of Lions (NW Mediterranean)." *Oceanologica Acta* vol. 23.
- 3 - Gaufrès P., Andres B., Dufois F., Interest and limits of oceanographic real time measurement on buoyancy beacon – Case of Rhône delta and gulf of Fos. Mediterranean Days of coastal and port engineering, Palermo 7-9 October 2008.