

ANARHICHADIDAE

atlantic wolffish

Anarhichas lupus

Linnaeus, 1758



oto. Adlava

HABITAT AND ECOLOGY

Demersal, mainly on rocky bottoms. Feeds on hard shelled organisms such as molluscs, crabs and echinoderms. Pelagic larvae settle to the substrate when 5-6 cm.

DISTRIBUTION

Mediterranean record: a single record from Varazze, Italy (Tortonese, 1958).

Original distribution: northern Atlantic Ocean from White Sea, North Sea to Cape Cod.

KEY REFERENCE

• Tortonese E. 1958. Primo ritrovamento di *Anarhichas lupus* L. (Piaces) nel Mediterraneo (Golfo di Genova). *Doriana*, 2(94): 1-4.

MURAENIDAE

moray eels

Gymnothorax reticularis

Bloch, 1795



Photo: N. Stern

HABITAT AND ECOLOGY

Inhabits rocky and open benthic soft substrate. Feeds on fish and large invertebrates. Eggs and larvae planktonic.

DISTRIBUTION

Mediterranean record: a single specimen was collected from Israel (Stern and Goren, 2013).

Original distribution: Indo-Pacific including the Red Sea.

KEY REFERENCE

• Stern N. and Goren M. 2013. First record of the moray eel *Gymnothorax reticularis*, Bloch, 1795 in the Mediterranean Sea, with a note on its taxonomy and distribution. *Zootaxa*, 3641(2): 197–200.